

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
9 June 2005 (09.06.2005)

PCT

(10) International Publication Number  
**WO 2005/051893 A2**

(51) International Patent Classification<sup>7</sup>: **C07C 235/00**

(21) International Application Number:  
PCT/US2004/037182

(22) International Filing Date:  
16 November 2004 (16.11.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/523,301 20 November 2003 (20.11.2003) US

GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **ELI LILLY AND COMPANY** [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **BUNEL, Emilio, Enrique** [CL/US]; 3277 Sunset Hills Boulevard, Thousand Oaks, CA 91362 (US). **GAJEWSKI, Robert, Peter** [US/US]; 1501 Friendship Drive, Indianapolis, IN 46217 (US). **JONES, Charles, David** [US/US]; 223 East Brunswick Avenue, Indianapolis, IN 46227 (US). **LU, Jianliang** [CN/US]; 11921 Castlestone Drive, Fishers, IN 46038 (US). **NAGPAL, Sunil** [US/US]; 5258 Comanche Trail, Carmel, IN 46033 (US). **MA, Tianwei** [CN/US]; 5676 Sapphire Drive, Carmel, IN 46033 (US). **YEE, Ying, Kwong** [US/US]; 5127 Briarstone Trace, Carmel, IN 46033 (US).

(74) Agents: **BOUDREAUX, William, R.** et al., **ELI LILLY AND COMPANY**, P. O. Box 6288, Indianapolis, IN 46206-6288 (US)

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,

**Declarations under Rule 4.17:**

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations* AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, *ARIPO patent* (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), *Eurasian patent* (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), *European patent* (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), *OAPI patent* (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- *of inventorship (Rule 4.17(iv)) for US only*

**Published:**

- *without international search report and to be republished upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: VITAMIN D RECEPTOR MODULATORS

(57) Abstract: The present invention relates to novel, non-secosteroidal, hydroxyl substituted, carbon-linked diaryl compounds with vitamin D receptor (VDR) modulating activity that are less hypercalcemic than 1,25 dihydroxy vitamin D<sub>3</sub>. These compounds are useful for treating bone disease and psoriasis.

WO 2005/051893 A2

-1-

## VITAMIN D RECEPTOR MODULATORS

### BACKGROUND OF THE INVENTION

5 Vitamin D<sub>3</sub> Receptor (VDR) is a ligand dependent transcription factor that belongs to the superfamily of nuclear hormone receptors. The VDR protein is 427 amino acids, with a molecular weight of ~50 kDa. The VDR ligand, 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> (the hormonally active form of Vitamin D) has its action mediated by its interaction with the nuclear receptor known as Vitamin D receptor ("VDR"). The VDR ligand, 1 $\alpha$ ,25-

10 dihydroxyvitamin D<sub>3</sub> (1 $\alpha$ ,25(OH)<sub>2</sub>D<sub>3</sub>) acts upon a wide variety of tissues and cells both related to and unrelated to calcium and phosphate homeostasis.

The activity 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> in various systems suggests wide clinical applications. However, use of conventional VDR ligands is hampered by their associated toxicity, namely hypercalcemia (elevated serum calcium). Currently, 1 $\alpha$ ,25(OH)<sub>2</sub>D<sub>3</sub>,

15 marketed as Rocaltrol® pharmaceutical agent ( product of Hoffmann-La Roche), is administered to kidney failure patients undergoing chronic kidney dialysis to treat hypocalcemia and the resultant metabolic bone disease. Other therapeutic agents, such as Calcipotriol® (synthetic analog of 1 $\alpha$ ,25(OH)<sub>2</sub>D<sub>3</sub> ) show increased separation of binding affinity on VDR from hypercalcemic activity.

20 Chemical modifications of 1 $\alpha$ ,25(OH)<sub>2</sub>D<sub>3</sub> have yielded analogs with attenuated calcium mobilization effects (R. Bouillon et. al., Endocrine Rev. 1995, 16, 200-257). One such analog, Dovonex ® pharmaceutical agent (product of Bristol-Meyers Squibb Co.), is currently used in Europe and the United States as a topical treatment for mild to moderate psoriasis (K. Kragballe et. al., Br. J. Dermatol. 1988, 119, 223-230).

25 Other Vitamin D<sub>3</sub> mimics have been described in the publication, Vitamin D Analogs: Mechanism of Action of Therapeutic Applications, by Nagpal, S.; Lu, J.; Boehm, M. F., Curr. Med. Chem. 2001, 8, 1661-1679.

Although some degree of separation between the beneficial action and calcium raising (calcemic) effects has been achieved with these VDR ligands, to date the

30 separation has been insufficient to allow for oral administration to treat conditions such as osteoporosis, cancers, leukemias, and severe psoriasis.

-2-

One example of a major class of disorder that could benefit from VDR mediated biological efficacy in the absence of hypercalcemia is osteoporosis. Osteoporosis is a systemic disorder characterized by decreased bone mass and microarchitectural deterioration of bone tissue leading to bone fragility and increased susceptibility to fractures of the hip, spine, and wrist (World Health Organization WHO 1994). Osteoporosis affects an estimated 75 million people in the United States, Europe, and Japan.

Within the past few years, several antiresorptive therapies have been introduced. These include bisphosphonates, hormone replacement therapy (HRT), a selective estrogen receptor modulator (SERM), and calcitonins. These treatments reduce bone resorption, bone formation, and increase bone density. However, none of these treatments increase true bone volume nor can they restore lost bone architecture.

Another major disorder that could benefit from VDR mediated biological activity is psoriasis. Psoriasis is one of the most common dermatologic diseases and is a chronic inflammatory skin condition characterized by erythematous, sharply demarcated papules and rounded plaques, covered by silvery micaceous scale.

Synthetic VDR ligands with reduced calcemic potential have been synthesized. For example, a class of bis-phenyl compounds stated to mimic  $1\alpha, 25$ -dihydroxyvitamin  $D_3$  is described in US Patent No. 6,218,430 and the article; "Novel nonsecosteroidal vitamin D mimics exert VDR-modulating activities with less calcium mobilization than  $1\alpha, 25$ -Dihydroxyvitamin  $D_3$ " by Marcus F. Boehm, et. al., Chemistry & Biology 1999, Vol 6, No. 5, pgs. 265-275.

Synthetic VDR ligands having an aryl-thiophene nucleus are described in United States provisional patent application SN 60/384151, filed 29 May 2002.


There remains a need for improved treatments using alternative or improved pharmaceutical agents that mimic  $1\alpha, 25$ -dihydroxyvitamin  $D_3$  to stimulate bone formation, restore bone quality, and treat other diseases without the attendant disadvantage of hypercalcemia.

## SUMMARY OF THE INVENTION


Novel compounds having a nucleus of formula "(A)" have been found effective as

-3-

Vitamin D Receptor (VDR) modulators:



The compounds of the invention with VDR modulating activities are represented by  
 5 formula (I)



wherein the variables R, R', R<sub>PH</sub>, R<sub>1</sub>, R<sub>2</sub>, L<sub>1</sub>, L<sub>2</sub>, L<sub>3</sub>, R<sub>BOH</sub>, and R<sub>C</sub> are as hereinafter defined. It is a discovery of this invention that compounds described herein display the  
 10 desirable cell differentiation and antiproliferative effects of 1,25(OH)<sub>2</sub>D<sub>3</sub> with reduced calcium mobilization (calcemic) effects if substituent R<sub>BOH</sub> bears a hydroxyl group and substituent R<sub>C</sub> possesses a carbon atom linked group directly bonded (i.e., with no intervening non-carbon atom) to the aryl nucleus.

In another aspect, the present invention is directed towards pharmaceutical  
 15 compositions containing pharmaceutically effective amounts of compounds of formulae (I) or a pharmaceutically acceptable salt or prodrug thereof, either singly or in combination, together with pharmaceutically acceptable carriers and/or auxiliary agents.

Another aspect of the invention is a pharmaceutical formulation for treatment or prevention of osteoporosis containing pharmaceutically effective amounts of the vitamin  
 20 D receptor modulator compound of formula (I) alone or together with pharmaceutically effective amounts of co-agents conventionally used for the treatment of osteoporosis.

Another aspect of the invention is a pharmaceutical formulation for treatment or prevention of psoriasis containing pharmaceutically effective amounts of the vitamin D  
 25 receptor modulator compound of formula (I) alone or together with pharmaceutically effective amounts of co-agents conventionally used for the treatment of psoriasis.

Another aspect of the invention is a pharmaceutical formulation for treatment or prevention of prostate cancer containing pharmaceutically effective amounts of the

-4-

vitamin D receptor modulator compound of formula (I) alone or together with pharmaceutically effective amounts of co-agents conventionally used for the treatment of prostate cancer.

Another aspect of the invention is to use the compounds of the invention to treat disease states responsive to Vitamin D receptor ligands.

Another aspect of the invention is the prevention and treatment of acne, actinic keratosis, alopecia, Alzheimer's disease, autoimmune induced diabetes, bone fracture healing, breast cancer, Crohn's disease, prostate cancer, benign prostatic hyperplasia, bladder cancer, colon cancer, Type I diabetes, host-graft rejection, hypercalcemia, Type II diabetes, leukemia, multiple sclerosis, insufficient sebum secretion, osteomalacia, osteoporosis, insufficient dermal firmness, insufficient dermal hydration, myelodysplastic syndrome, psoriatic arthritis, psoriasis, renal osteodystrophy, rheumatoid arthritis, scleroderma, seborrheic dermatitis, skin cancer, systemic lupus erythematosus, skin cell protection from Mustard vesicants, ulcerative colitis, and wrinkles; by administering to a mammal in need thereof a pharmaceutically effective amount of a compound of Formula I.

#### DETAILED DESCRIPTION OF THE INVENTION

##### Definitions:

The term, "abscess" refers to adverse complications often associated with surgery, trauma, or diseases that predispose the host to abscess formation from encapsulated bacteria lymphocytes, macrophages, and etc.

The term, "adhesion" refers to the adverse and abnormal union of surfaces normally separate by the formation of new fibrous tissue resulting from an inflammatory process.

The term, "compound of the invention" refers to a compound represented by any of (i) formulae I, II, III and its preferred embodiments, (ii) the product of any example set out herein, or (iii) a compound identified in the species coded AA-1 to AA-33, BB-1 to BB-33, CC-1 to CC-44, or a salt or prodrug derivative of (i), (ii), or (iii).

The term, "Active Ingredient" means a compound of the invention.

The term, "Mustard" is inclusive of both sulfur mustards and nitrogen mustards, either alone or in any combination. Exemplary of such compounds are the vesicants;

-5-






- bis(2-chloroethyl) sulfide (Chemical Agent Symbol HD),  $\text{Cl}(\text{CH}_2)_2\text{S}(\text{CH}_2)_2\text{Cl}$  1,2-bis(2-chloroethylthio)ethane (Chemical Agent Symbol Q),  $\text{Cl}(\text{CH}_2)_2\text{S}(\text{CH}_2)_2\text{S}(\text{CH}_2)_2\text{Cl}$ ; bis(2-chloroethylthioethyl) ether,  $\text{Cl}(\text{CH}_2)_2\text{S}(\text{CH}_2)\text{O}(\text{CH}_2)_2\text{S}(\text{CH}_2)_2\text{Cl}$  (Chemical Agent Symbol T); tris(2-chloroethyl) amine (Chemical Agent Symbol HN3)  $\text{N}(\text{CH}_2\text{CH}_2\text{Cl})_3$ ;
- 5 N-methyl-2,2'-dichlorodiethylamine (Chemical Agent Symbol NH2); and 2,2'-dichlorotriethylamine,  $\text{CH}_3\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2\text{Cl})_2$  (Chemical Agent Symbol NH1).

The term, “(Acidic Group)” means an organic group that acts as a proton donor capable of hydrogen bonding. Illustrative of an (Acidic Group) is a group selected from the following:

10

















-C(O)OH,

-5-tetrazolyl,




15

-6-




-7-

or



The term, "carbon atom linked group" is used to identify the chemical substituent  $R_C$  in the Formula I definition of compounds of the invention. Its defining characteristic is a carbon atom as the first atom and point of attachment to the aryl ring to which it is attached. For example in the structural formula (C):



the arrow identifies the carbon atom linked directly to the aryl nucleus of formula (I). All compounds of the invention contain a carbon atom linked group as the  $R_C$  substituent.

The term "alkenyl" refers to aliphatic groups wherein the point of attachment is a carbon-carbon double bond, for example vinyl, 1-propenyl, and 1-cyclohexenyl. Alkenyl groups may be straight-chain, branched-chain, cyclic, or combinations thereof, and may be optionally substituted. Suitable alkenyl groups have from 2 to about 20 carbon atoms.

The term "C<sub>1</sub>-C<sub>5</sub> alkyl" refers to saturated aliphatic groups including straight-chain, branched-chain, and cyclic groups and any combinations thereof. Examples of C<sub>1</sub>-C<sub>5</sub> alkyl groups are methyl; ethyl; n-propyl; 1-methylethyl; n-butyl; 1-methylpropyl; 2-methylpropyl; n-amyl; 1,1-dimethylpropyl; 1,2-dimethylpropyl; and 2,2-dimethylpropyl.

The term "cycloalkyl" includes organic radicals such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.


The term, "cycloalkenyl" includes organic radicals such as cyclopropenyl, cyclobutenyl, cyclopentenyl and cyclohexenyl.

The term, "C<sub>1</sub>-C<sub>5</sub> fluoroalkyl" is an alkyl group containing fluorine and includes organic radicals such as -CF<sub>3</sub>, -CHF<sub>2</sub>, -CH<sub>2</sub>F, -CF<sub>2</sub>CF<sub>3</sub>, -CHFCF<sub>3</sub>, -CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>CHF<sub>2</sub>, and -CH<sub>2</sub>CH<sub>2</sub>F, with -CF<sub>3</sub> being preferred.




-8-


The term, “-1,3-thiazolidine-2,4-dione-5-methylidene”, refers to the radical represented by the structural formula:




5 The term, “-CH<sub>2</sub>-C(O)-N-pyrrolidine” refers to the radical represented by the structural formula:




The term, “-CH<sub>2</sub>-N-pyrrolidin-2-one” refers to the radical represented by the structural formula:



10 The term, “-CH<sub>2</sub>-(1-methylpyrrolidin-2-one-3-yl)” refers to the organic radical represented by the structural formula:




The term, “1,3,4-oxadiazolin-2-one-5-yl” refers to the organic radical represented by the structural formula:




15

The term, “1,3,4-oxadiazolin-2-thione-5-yl” refers to the organic radical represented by the structural formula:

-9-




The term, "imidazolidine-2,4-dione-5-yl" refers to the organic radical represented by the structural formula:



5

The term, "isoxazol-3-ol-5-yl" refers to the organic radical represented by the structural formula:




The dotted line symbol crossing a solid line representing a bond



10


The univalent symbol "-O" in any structural formula is a hydroxyl group (-OH). means that the bond so marked is the bond of attachment, for example, the group;



is attached to a phenyl ring of the parent diaryl nucleus to provide a compound of the invention as shown;

15

-10-



The term, "mammal" includes humans.

The term "halo" refer to fluorine, chlorine, bromine, and iodine.

5 The term, "terminal hydroxyalkyl" is a group selected from

3-methyl-3-hydroxypentyl,

3-methyl-3-hydroxypentenyl,

3-methyl-3-hydroxypentynyl,

3-ethyl-3-hydroxypentyl,

10 3-ethyl-3-hydroxypentenyl,

3-ethyl-3-hydroxypentynyl,

3-ethyl-3-hydroxy-4-methylpentyl,

3-ethyl-3-hydroxy-4-methylpentenyl,

3-ethyl-3-hydroxy-4-methylpentynyl,

15 3-propyl-3-hydroxypentyl,

3-propyl-3-hydroxypentenyl,


3-propyl-3-hydroxypentynyl,

1-hydroxy-2-methyl-1-(methylethyl)propyl,

1-hydroxycycloalkenyl; and


20 1-hydroxycycloalkyl.

The term, "3-methyl-3-hydroxypentyl" refers to the radical having the structural formula:




25 The term, "3-methyl-3-hydroxypentenyl" refers to the radical having the structural formula (both cis and trans isomers):


-11-




The term, "3-methyl-3-hydroxypentynyl" refers to the radical having the structural formula:



5 The term, "3-ethyl-3-hydroxypentyl" refers to the radical having the structural formula:




The term, "3-ethyl-3-hydroxypentenyl" refers to the radical having the structural formula (both cis and trans isomers):




10

The term, "3-ethyl-3-hydroxypentynyl" refers to the radical having the structural formula:




15 The term, "3-propyl-3-hydroxypentyl" refers to the radical having the structural formula:




The term, "3-propyl-3-hydroxypentenyl" refers to the radical having the structural formula (both cis and trans isomers):

-12-




The term, "3-propyl-3-hydroxypentynyl" refers to the radical having the structural formula:




5


The term, "3-ethyl-3-hydroxy-4-methylpentyl" refers to the radical having the structural formula:




The term, "3-ethyl-3-hydroxy-4-methylpentenyl" refers to the radical having the structural formula (both cis and trans isomers):



The term, "3-ethyl-3-hydroxy-4-methylpentynyl" refers to the radical having the structural formula:




The term, "1-hydroxy-2-methyl-1-(methylethyl)propyl" refers to the radical having the structural formula:



-13-


The term, "1-hydroxycycloalkenyl" refers to a radical selected from 1-hydroxycyclopentenyl, 1-hydroxycyclohexenyl, 1-hydroxycycloheptenyl, or 1-hydroxycyclooctenyl.

5 The term "hydroxycycloalkyl" refers to a radical having the general structural formula:



where  $w$  is an integer from 1 to 6 and the hydroxyl radical is substituted on any ring carbon atom.

10 The term "1-hydroxycycloalkyl" refers to a radical having the general structural formula:



Examples of 1-hydroxycycloalkyl radicals are

1-hydroxycyclopropyl, 1-hydroxycyclobutyl, 1-hydroxycyclopentyl,  
1-hydroxycyclohexyl, 1-hydroxycycloheptyl, and 1-hydroxycyclooctyl.

15 The abbreviation, "Me" means methyl.

The abbreviation, "Et" means ethyl.

The abbreviation, "iPr" means 1-methylethyl.

The abbreviation, "nPr" means n-propyl.

The abbreviation, "3Me3OH-Pentyl" means 3-methyl-3-hydroxypentyl.

20 The abbreviation, "3Me3OH-Pentenyl" means 3-methyl-3-hydroxypentenyl

The abbreviation, "3Me3OH-Pentynyl" means 3-methyl-3-hydroxypentynyl

The abbreviation, "3Et3OH-Pentyl" means 3-ethyl-3-hydroxypentyl.

The abbreviation, "3Et3OH-Pentenyl" means 3-ethyl-3-hydroxypentenyl

The abbreviation, "3Et3OH-Pentynyl" means 3-ethyl-3-hydroxypentynyl

25 The abbreviation, "3Pr3OH-Pentyl" means 3-propyl-3-hydroxypentyl.

The abbreviation, "3Pr3OH-Pentenyl" means 3-propyl-3-hydroxypentenyl.

-14-

The abbreviation, "3Pr3OH-Pentynyl" means 3-propyl-3-hydroxypentynyl.

The abbreviation, "3Et3OH4Me-Pentynyl" means 3-ethyl-3-hydroxy-4-methylpentynyl.


The abbreviation, "3Et3OH4Me-Pentenyl" means 3-ethyl-3-hydroxy-4-methylpentenyl,

5 The abbreviation, "3Et3OH4Me-Pentynyl" means 3-ethyl-3-hydroxy-4-methylpentynyl.

The abbreviation, "1OH2Me1MeEt-Propyl" means 1-hydroxy-2-methyl-1-(methylethyl)propyl.

# 10 Compounds of the Invention:

The compounds of the invention with vitamin receptor modulating (VDRM) activities are represented by formula (I) or a pharmaceutically acceptable salt or a prodrug derivative thereof:



15

wherein;

R and R' are independently C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>1</sub>-C<sub>5</sub> fluoroalkyl, or together R and R' form a substituted or unsubstituted, saturated or unsaturated carbocyclic ring having from  
20 3 to 8 carbon atoms;






R<sub>PH</sub> is hydrogen or methyl;

R<sub>1</sub> and R<sub>2</sub> are independently selected from the group consisting of hydrogen, halo, C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>1</sub>-C<sub>5</sub> fluoroalkyl, -O-C<sub>1</sub>-C<sub>5</sub> alkyl, -S-C<sub>1</sub>-C<sub>5</sub> alkyl, -O-C<sub>1</sub>-C<sub>5</sub> fluoroalkyl, -CN, -NO<sub>2</sub>, acetyl, -S-C<sub>1</sub>-C<sub>5</sub> fluoroalkyl, C<sub>2</sub>-C<sub>5</sub> alkenyl, C<sub>3</sub>-C<sub>5</sub> cycloalkyl,  
25 and C<sub>3</sub>-C<sub>5</sub> cycloalkenyl;

L<sub>1</sub> and L<sub>2</sub> and L<sub>3</sub> are independently divalent linking groups independently selected from the group consisting of


-15-

a bond ,






R40

R40








R40





-16-



- 5 where m is 0, 1, or 2; X<sub>1</sub> is oxygen or sulfur, and each R<sub>40</sub> is independently hydrogen, C<sub>1</sub>-C<sub>5</sub> alkyl, or C<sub>1</sub>-C<sub>5</sub> fluoroalkyl;

R<sub>BOH</sub> is

- 3-methyl-3-hydroxypentyl,  
 3-methyl-3-hydroxypentenyl,  
 10 3-methyl-3-hydroxypentynyl,  
 3-ethyl-3-hydroxypentyl,  
 3-ethyl-3-hydroxypentenyl,  
 3-ethyl-3-hydroxypentynyl,  
 3-ethyl-3-hydroxy-4-methylpentyl,  
 15 3-ethyl-3-hydroxy-4-methylpentenyl,  
 3-ethyl-3-hydroxy-4-methylpentynyl,  
 3-propyl-3-hydroxypentyl,

-17-

5 3-propyl-3-hydroxypentenyl,  
3-propyl-3-hydroxypentynyl,  
1-hydroxy-2-methyl-1-(methylethyl)propyl  
1-hydroxycyclopentenyl,  
1-hydroxycyclohexenyl,  
1-hydroxycycloheptenyl,  
1-hydroxycyclooctenyl,  
1-hydroxycyclopropyl,  
1-hydroxycyclobutyl,  
10 1-hydroxycyclopentyl,  
1-hydroxycyclohexyl,  
1-hydroxycycloheptyl, or  
1-hydroxycyclooctyl;

provided that when

15  $R_{BOH}$  is

3-methyl-3-hydroxypentyl,  
3-methyl-3-hydroxypentenyl,  
3-methyl-3-hydroxypentynyl,  
3-ethyl-3-hydroxypentyl,  
20 3-ethyl-3-hydroxypentenyl,  
3-ethyl-3-hydroxypentynyl,  
3-ethyl-3-hydroxy-4-methylpentyl,  
3-ethyl-3-hydroxy-4-methylpentenyl,  
3-ethyl-3-hydroxy-4-methylpentynyl,  
25 3-propyl-3-hydroxypentyl,  
3-propyl-3-hydroxypentenyl,  
3-propyl-3-hydroxypentynyl, or  
1-hydroxy-2-methyl-1-(methylethyl)propyl;

then  $L_1$  and  $L_2$  combine as a bond; and

-18-

 $R_C$  is

- 5
- 10
- 15
- 20
- 25
- CO<sub>2</sub>H,
  - CO<sub>2</sub>Me,
  - CO<sub>2</sub>Et,
  - C(O)CH<sub>2</sub>S(O)Me,
  - C(O)CH<sub>2</sub>S(O)Et,
  - C(O)CH<sub>2</sub>S(O)<sub>2</sub>Me,
  - C(O)CH<sub>2</sub>S(O)<sub>2</sub>Et,
  - C(O)CH<sub>2</sub>CH<sub>2</sub>S(O)Me,
  - C(O)CH<sub>2</sub>CH<sub>2</sub>S(O)Et,
  - C(O)CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>Me,
  - C(O)CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>Et,
  - C(O)CHMeCH<sub>2</sub>CO<sub>2</sub>H
  - C(O)C(O)OH,
  - C(O)C(O)NH<sub>2</sub>,
  - C(O)C(O)NHMe,
  - C(O)C(O)NMe<sub>2</sub>,
  - C(O)NH<sub>2</sub>, C(O)NMe<sub>2</sub>,
  - C(O)NHS(O)Me,
  - C(O)NHSO<sub>2</sub>Me,
  - C(O)-NH-5-tetrazolyl,
  - C(O)NMe-5-tetrazolyl,
  - C(O)NHS(O)Me,
  - C(O)NHS(O)Et,
  - C(O)NHSO<sub>2</sub>Me,
  - C(O)NHSO<sub>2</sub>Et,

-19-

- 5
- C(O)NHS(O)iPr,  
-C(O)NHSO<sub>2</sub>iPr,  
-C(O)NHS(O)nPr,  
-C(O)NHSO<sub>2</sub>nPr,  
-C(O)NHCH<sub>2</sub>S(O)Me,  
-C(O)NHCH<sub>2</sub>S(O)Et,  
-C(O)NHCH<sub>2</sub>SO<sub>2</sub>Me,  
-C(O)NHCH<sub>2</sub>SO<sub>2</sub>Et,  
10 -C(O)NHCH<sub>2</sub>CH<sub>2</sub>S(O)Me,  
-C(O)NHCH<sub>2</sub>CH<sub>2</sub>S(O)Et,  
-C(O)NHCH<sub>2</sub>CH<sub>2</sub>SO<sub>2</sub>Me,  
-C(O)NHCH<sub>2</sub>CH<sub>2</sub>SO<sub>2</sub>Et,  
-C(O)NH<sub>2</sub>,  
-C(O)NMe<sub>2</sub>,  
15 -C(O)NH-CH<sub>2</sub>-C(O)OH,  
-C(O)NH-CH(Me)-C(O)OH,  
-C(O)NH-CH(F)-C(O)OH,  
-C(O)NH-CH(CF<sub>3</sub>)-C(O)OH,  
-C(O)NH-CH(OH)-C(O)OH,  
20 -C(O)NH-CH(cyclopropyl)-C(O)OH,  
-C(O)NH-C(Me)<sub>2</sub>-C(O)OH,  
-C(O)NH-C(Me)<sub>2</sub>-C(O)OH,  
-C(O)NH-CF(Me)-C(O)OH,  
-C(O)NH-C(Me)(CF<sub>3</sub>)-C(O)OH,  
25 -C(O)NH-C(Me)(OH)-C(O)OH,  
-C(O)NH-C(Me)(cyclopropyl)-C(O)OH,  
-C(O)NMe-CH<sub>2</sub>-C(O)OH,  
-C(O)NMe-CH(Me)-C(O)OH,  
-C(O)NMe-CH(F)-C(O)OH,  
30 -C(O)NMe-CH(CF<sub>3</sub>)-C(O)OH,  
-C(O)NMe-CH(OH)-C(O)OH,  
-C(O)NMe-CH(cyclopropyl)-C(O)OH,

-20-

- 5
- C(O)NMe-C(Me)<sub>2</sub>-C(O)OH,  
-C(O)NMe-CF(Me)-C(O)OH,  
-C(O)NMe-C(Me)(CF<sub>3</sub>)-C(O)OH,  
-C(O)NMe-C(Me)(OH)-C(O)OH,  
-C(O)NMe-C(Me)(cyclopropyl)-C(O)OH,  
-CH<sub>2</sub>-CO<sub>2</sub>H,  
-CH<sub>2</sub>-5-tetrazolyl,  
-CH<sub>2</sub>CO<sub>2</sub>Me,  
-CH<sub>2</sub>CO<sub>2</sub>Et,  
10 -CH<sub>2</sub>NHS(O)Me,  
-CH<sub>2</sub>NHS(O)Et,  
-CH<sub>2</sub>NHSO<sub>2</sub>Me,  
-CH<sub>2</sub>NHSO<sub>2</sub>Et,  
-CH<sub>2</sub>NHS(O)iPr,  
15 -CH<sub>2</sub>NHSO<sub>2</sub>iPr,  
-CH<sub>2</sub>NHS(O)nPr,  
-CH<sub>2</sub>NHSO<sub>2</sub>nPr,  
-CH<sub>2</sub>NHCH<sub>2</sub>CH<sub>2</sub>SO<sub>2</sub>CH<sub>3</sub>,  
-CH<sub>2</sub>NH(CH<sub>2</sub>CO<sub>2</sub>H),  
20 -CH<sub>2</sub>N(C(O)Me)(CH<sub>2</sub>CO<sub>2</sub>H),  
-CH<sub>2</sub>-N-pyrrolidin-2-one,  
-CH<sub>2</sub>-(1-methylpyrrolidin-2-one-3-yl),  
-CH<sub>2</sub>S(O)Me,  
-CH<sub>2</sub>S(O)Et,  
25 -CH<sub>2</sub>S(O)<sub>2</sub>Me,  
-CH<sub>2</sub>S(O)<sub>2</sub>Et,  
-CH<sub>2</sub>S(O)iPr,  
-CH<sub>2</sub>S(O)<sub>2</sub>iPr,  
-CH<sub>2</sub>S(O)nPr,  
30 -CH<sub>2</sub>S(O)<sub>2</sub>nPr,  
-CH<sub>2</sub>CO<sub>2</sub>H, CH<sub>2</sub>C(O)NH<sub>2</sub>,  
-CH<sub>2</sub>C(O)NMe<sub>2</sub>,

-21-

- 5                   -CH<sub>2</sub>C(O)NHMe,  
                  -CH<sub>2</sub>C(O)-N-pyrrolidine,  
                  -CH<sub>2</sub>S(O)<sub>2</sub>Me,  
                  -CH<sub>2</sub>S(O)Me,  
                  -CH(OH) CO<sub>2</sub>H,  
                  -CH(OH)C(O)NH<sub>2</sub>,  
                  -CH(OH)C(O)NHMe,  
                  -CH(OH)C(O)NMe<sub>2</sub>,  
                  -CH(OH)C(O)NEt<sub>2</sub>,  
10               -CH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub>H,  
                  -CH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub>Me,  
                  -CH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub>Et,  
                  -CH<sub>2</sub>CH<sub>2</sub>C(O)NH<sub>2</sub>,  
                  -CH<sub>2</sub>CH<sub>2</sub>C(O)NHMe,  
15               -CH<sub>2</sub>CH<sub>2</sub>C(O)NMe<sub>2</sub>,  
                  -CH<sub>2</sub>CH<sub>2</sub>-5-tetrazolyl,  
                  -CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>Me,  
                  -CH<sub>2</sub>CH<sub>2</sub>S(O)Me,  
                  -CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>Et,  
20               -CH<sub>2</sub>CH<sub>2</sub>S(O) Et,  
                  -CH<sub>2</sub>CH<sub>2</sub>S(O)iPr,  
                  -CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>iPr,  
                  -CH<sub>2</sub>CH<sub>2</sub>S(O)nPr,  
                  -CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>nPr,  
25               -CH<sub>2</sub>CH<sub>2</sub>S(O)NH<sub>2</sub>,  
                  -CH<sub>2</sub>CH<sub>2</sub>S(O)NHMe,  
                  -CH<sub>2</sub>CH<sub>2</sub>S(O)NMe<sub>2</sub>,  
                  -CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>NH<sub>2</sub>,  
                  -CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>NHMe,  
30               -CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>NMe<sub>2</sub>,  
                  -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>S(O)Me,  
                  -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>S(O)Et,


-22-

-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>Me,-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>Et,-CH(Me)CH<sub>2</sub>C(O)OH,-C(Me)<sub>2</sub>CH<sub>2</sub>C(O)OH,









5

-SO<sub>3</sub>H,





-5-tetrazolyl,




10



-23-



or



-1,3,4-oxadiazolin-2-one-5-yl,

-imidazolidine-2,4-dione-5-yl,

-1,3-thiazolidine-2,4-dione-5-methylidene,

-isoxazol-3-ol-yl, or

-1,3,4-oxadiazolin-2-thione-5-yl.


5

The combination of  $L_1$  and  $L_2$  as a bond (divalent linking group) is understood to mean




-24-

that group  $R_{BOH}$ , is attached directly to the phenyl nucleus, for example, as in the following structural formula:



- 5 where  $R_{BOH}$  is 3-ethyl-3-hydroxypentenyl.

Preferred compounds of the invention with VDR modulating activities are represented by formula (II) or a pharmaceutically acceptable salt or an ester prodrug derivative thereof:



10




wherein;

R and R' are independently methyl or ethyl;


R1 and R2 are independently hydrogen, halo,  $-CF_3$ , methyl, ethyl, or cyclopropyl;

$L_1$  and  $L_2$  are independently divalent linking groups independently selected from

a bond ,



-25-



5 where m is 0 or 1;

R<sub>BOH</sub> is selected from

1-hydroxycyclopentenyl,  
 1-hydroxycyclohexenyl,  
 1-hydroxycyclopentyl, or  
 10 1-hydroxycyclohexyl, and


Z<sub>C</sub> is a group selected from

-CO<sub>2</sub>H,  
 -CO<sub>2</sub>Me,  
 -CO<sub>2</sub>Et,  
 15 -C(O)NH<sub>2</sub>,  
 -C(O)NMe<sub>2</sub>,  
 -C(O)NH-CH<sub>2</sub>-C(O)OH,  
 -C(O)NH-CH(Me)-C(O)OH,  
 -C(O)NH-CH(F)-C(O)OH,  
 20 -C(O)NH-CH(CF<sub>3</sub>)-C(O)OH,  
 -C(O)NH-CH(OH)-C(O)OH,  
 -C(O)NH-CH(cyclopropyl)-C(O)OH,  
 -C(O)NH-C(Me)<sub>2</sub>-C(O)OH,  
 -C(O)NH-C(Me)<sub>2</sub>-C(O)OH,  
 25 -C(O)NH-CF(Me)-C(O)OH,  
 -C(O)NH-C(Me)(CF<sub>3</sub>)-C(O)OH,  
 -C(O)NH-C(Me)(OH)-C(O)OH,  
 -C(O)NH-C(Me)(cyclopropyl)-C(O)OH,  
 -C(O)NMe-CH<sub>2</sub>-C(O)OH,  
 30 -C(O)NMe-CH(Me)-C(O)OH,  
 -C(O)NMe-CH(F)-C(O)OH,

-26-

-C(O)NMe-CH(CF<sub>3</sub>)-C(O)OH,  
 -C(O)NMe-CH(OH)-C(O)OH,  
 -C(O)NMe-CH(cyclopropyl)-C(O)OH,  
 -C(O)NMe-C(Me)<sub>2</sub>-C(O)OH,  
 -C(O)NMe-CF(Me)-C(O)OH,  
 -C(O)NMe-C(Me)(CF<sub>3</sub>)-C(O)OH,  
 -C(O)NMe-C(Me)(OH)-C(O)OH,  
 -C(O)NMe-5-tetrazolyl,  
 -C(O)NMe-C(Me)(cyclopropyl)-C(O)OH, or  
 -C(O)-NH-5-tetrazolyl.

Preferred compounds of the invention with VDR modulating activities are represented by formula (III) or a pharmaceutically acceptable salt or an ester prodrug derivative thereof:



wherein;

R and R' are independently methyl or ethyl;

R<sub>1</sub> and R<sub>2</sub> are independently hydrogen, halo, -CF<sub>3</sub>, methyl, ethyl, or cyclopropyl;

R<sub>BOH</sub> is selected from

3-methyl-3-hydroxypentyl,  
 3-methyl-3-hydroxypentenyl,  
 3-methyl-3-hydroxypentynyl,  
 3-ethyl-3-hydroxypentyl,  
 3-ethyl-3-hydroxypentenyl,  
 3-ethyl-3-hydroxypentynyl,  
 3-propyl-3-hydroxypentyl,  
 3-propyl-3-hydroxypentynyl,  
 3-propyl-3-hydroxypentenyl,  
 3-ethyl-3-hydroxy-4-methylpentyl,

-27-

3-ethyl-3-hydroxy-4-methylpentenyl,  
3-ethyl-3-hydroxy-4-methylpentynyl, or  
1-hydroxy-2-methyl-1-(methylethyl)propyl; and

Z<sub>C</sub> is a group selected from

- 5                   -CO<sub>2</sub>H,  
                  -CO<sub>2</sub>Me,  
                  -CO<sub>2</sub>Et,  
                  -C(O)NH<sub>2</sub>,  
                  -C(O)NMe<sub>2</sub>,  
10               -C(O)NH-CH<sub>2</sub>-C(O)OH,  
                  -C(O)NH-CH(Me)-C(O)OH,  
                  -C(O)NH-CH(F)-C(O)OH,  
                  -C(O)NH-CH(CF<sub>3</sub>)-C(O)OH,  
                  -C(O)NH-CH(OH)-C(O)OH,  
15               -C(O)NH-CH(cyclopropyl)-C(O)OH,  
                  -C(O)NH-C(Me)<sub>2</sub>-C(O)OH,  
                  -C(O)NH-C(Me)<sub>2</sub>-C(O)OH,  
                  -C(O)NH-CF(Me)-C(O)OH,  
                  -C(O)NH-C(Me)(CF<sub>3</sub>)-C(O)OH,  
20               -C(O)NH-C(Me)(OH)-C(O)OH,  
                  -C(O)NH-C(Me)(cyclopropyl)-C(O)OH,  
                  -C(O)NMe-CH<sub>2</sub>-C(O)OH,  
                  -C(O)NMe-CH(Me)-C(O)OH,  
                  -C(O)NMe-CH(F)-C(O)OH,  
25               -C(O)NMe-CH(CF<sub>3</sub>)-C(O)OH,  
                  -C(O)NMe-CH(OH)-C(O)OH,  
                  -C(O)NMe-CH(cyclopropyl)-C(O)OH,  
                  -C(O)NMe-C(Me)<sub>2</sub>-C(O)OH,  
                  -C(O)NMe-CF(Me)-C(O)OH,  
30               -C(O)NMe-C(Me)(CF<sub>3</sub>)-C(O)OH,  
                  -C(O)NMe-C(Me)(OH)-C(O)OH,  
                  -C(O)NMe-5-tetrazolyl,

-28-


-C(O)NMe-C(Me)(cyclopropyl)-C(O)OH, or  
-C(O)-NH-5-tetrazolyl.

Particularly preferred salts of the compounds Formulae I and II and III are sodium  
5 and potassium.


Particularly preferred ester prodrugs of the compounds of Formulae I and II and III  
are the methyl ester, ethyl ester, N,N-diethylglycolamido ester, and the morpholinylethyl  
ester.

Particularly preferred are compounds or a pharmaceutically acceptable salt or  
10 prodrug derivative thereof selected from (AA-1) to (AA-33) and mixtures thereof, as  
follows:

AA-1)




AA-2)




AA-3)


-29-




AA-4)



AA-5)




AA-6)




AA-7)


-30-




AA-8)



AA-9)




AA-10)




AA-11)


-31-




AA-12)



AA-13)




AA-14)




AA-15)




-32-




AA-16)



AA-17)




AA-18)




AA-19)


-33-




AA-20)



AA-21)




AA-22)




AA-23)


-34-




AA-24)



AA-25)




AA-26)




AA-27)

-35-




AA-28)




AA-29)

5




AA-30)




AA-31)


-36-



AA-32)



AA-33)




5

Additional particularly preferred are compounds or a pharmaceutically acceptable salt or prodrug derivative thereof selected from (BB-1) to (BB-33) and mixtures thereof, as follows:


10

BB-1)


-37-




BB-2)



BB-3)




BB-4)




BB-5)


-38-




BB-6)



BB-7)




BB-8)




BB-9)


-39-




BB-10)



BB-11)




BB-12)




BB-13)




-40-




BB-14)



BB-15)




BB-16)




BB-17)


-41-




BB-18)



BB-19)




BB-20)




BB-21)


-42-




BB-22)



BB-23)




BB-24)




BB-25)


-43-




BB-26)



BB-27)




BB-28)




BB-29)


-44-




BB-30)



BB-31)




BB-32)




BB-33)

-45-




Additional particularly preferred are compounds or a pharmaceutically acceptable salt or prodrug derivative thereof selected from (CC-1) to (CC-44) and mixtures thereof,  
 5 as follows:

CC-1)




CC-2)




10

CC-3)


-46-




CC-4)



CC-5)




CC-6)




CC-7)


-47-




CC-8)



CC-9)




CC-10)




CC-11)




-48-




CC-12)



CC-13)




CC-14)




CC-15)

-49-




CC-16)




5

CC-17)




CC-18)




10

CC-19)


-50-




CC-20)



CC-21)




CC-22)




CC-23)


-51-




CC-24)



CC-25)




CC-26)




CC-27)


-52-



CC-28)




CC-29)




5

CC-30)




CC-31)


-53-




CC-32)



CC-33)




CC-34)




CC-35)

-54-




CC-36)




CC-37)

5




CC-38)




CC-39)


-55-




CC-40)



CC-41)





CC-42)




CC-43)




-56-



5




Particularly preferred compounds of the invention are



10

or




-57-

or pharmaceutically acceptable salts or prodrug derivatives thereof.


Scheme 1: Synthesis of Unsymmetrical  
Central Link Diphenyl Scaffold

5



-58-

Scheme 2: Synthesis of Trans-Pentenol  
Phenyl Alkyl Phenyl Amide-Acids




-59-

## Scheme 3: Synthesis of Pentynol


## Phenyl Alkyl Phenyl Acids

5



-60-

Scheme 4: Synthesis of Cis-Pentenol  
Phenyl Alkyl Phenyl Amide-Acids



5

Preparation of unsymmetrical central link diphenyl scaffold (Scheme 1).

A 3-Substituted-4-hydroxybenzoic acid (1a) is coupled with EDCI/N-methoxy-N-methoxyamine/DMAP and alkylated with benzyl bromide to give amide 2. Amide 2 is sequentially reacted with R<sub>2</sub>MgBr and R<sub>3</sub>MgBr Grignard reagents to afford tertiary alcohol 3. Alcohol 3 is reacted with 2-substituted phenol 4 and BF<sub>3</sub>·OEt<sub>2</sub> to produce diphenylalkane 5. Diphenylalkane 5 is reacted with triflic anhydride/pyridine to give the triflate 6, which is methoxycarbonylated with Pd(OAc)<sub>2</sub>, (DPPF or DPPB), carbon monoxide, MeOH, and Et<sub>3</sub>N to give ester 7.

15

Synthesis of trans-Pentenol Phenyl Alkyl Phenyl Acids (Scheme 2).

Triflate 6 is sequentially reacted with 1) TMS-acetylene, PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>, Et<sub>3</sub>N, and DMF and 2) CsF and water to afford acetylene 8. Acetylene 8 is treated with Zn(OTf)<sub>2</sub>/t-butyl aldehyde/chiral auxiliary (with or without) to give alcohol 10. Alternatively, acetylene 8 is reacted with LiHMDS/ketone 9 to give alcohol 10. Alcohol 10 is reduced with LAH or

20

-61-

DiBAH to afford trans-pentenol 11. Trans-pentenol 11 is sequentially reacted with 1) Pd-C/H<sub>2</sub>; 2) Tf<sub>2</sub>O/pyridine; 3) Pd(OAc)<sub>2</sub>, DPPF, CO, MeOH, Et<sub>3</sub>N, DMF; 4) KOH/EtOH/H<sub>2</sub>O; 5) EDCI/Et<sub>3</sub>N/DMAP/ R<sub>4</sub>NHCH<sub>2</sub>CO<sub>2</sub>Me; and 6) LiOH/EtOH/H<sub>2</sub>O to give trans-pentenol amide-acid 12. For reaction step 3, DPPB and DMSO may be substituted for DPPF and DMF, respectively. Trans-pentenol amide-acid 12 is hydrogenated with Pd-C/H<sub>2</sub> afford amide-acid 19.

#### Synthesis of Pentynol Phenyl alkyl Phenyl Acids (Scheme 3).

Ester 7 is hydrogenolyzed with Pd-C/H<sub>2</sub> and reacted with Tf<sub>2</sub>O/pyridine to give triflate 13. Triflate 13 is sequentially reacted with 1) TMS-acetylene, PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>, Et<sub>3</sub>N, and DMF and 2) CsF and water to afford acetylene 14. Acetylene 14 is treated with Zn(OTf)<sub>2</sub>/t-butyl aldehyde/chiral auxiliary (with or without) to give alcohol 15. Alternatively, acetylene 14 is reacted with LiHMDS/ketone 9 to give alcohol 15. Alcohol 15 is hydrolyzed with KOH/EtOH/H<sub>2</sub>O to afford acid 16. Acid 16 is sequentially reacted with 1) EDCI/Et<sub>3</sub>N/DMAP/R<sub>4</sub>NHCH<sub>2</sub>CO<sub>2</sub>Me and 2) LiOH/EtOH/H<sub>2</sub>O to give amide-acid 17.

#### Synthesis of Cis-Pentenol Phenyl alkyl Phenyl Acids (Scheme 4).

Amide-acid 17 is hydrogenated with Lindlar catalyst to afford cis-pentenol amide-acid 18. Cis-pentenol amide-acid 18 is hydrogenated with Pd-C/H<sub>2</sub> afford amide-acid 19. Alternatively, amide-acid 17 is hydrogenated with Pd-C/H<sub>2</sub> afford amide-acid 19.

### EXAMPLES

#### General Experimental Conditions:

The starting material/intermediate is the compound from the immediate preceding experimental unless otherwise indicated.

All reactions are performed under nitrogen/argon atmosphere, in a stirred reaction vessel, and at room temperature unless indicated otherwise.

Unless otherwise indicated, the organic layer is MgSO<sub>4</sub>/Na<sub>2</sub>SO<sub>4</sub> dried is defined as stirring the solution with a dessicant for 5-15 m and filtering off the dessicant to give an anhydrous filtrate.

-62-

For analogous multi-step reaction procedures, the yield is given either for the ultimate step or overall multi-steps as indicated.

Solutions are "concentrated" at a range of 25-75 °C with reduced pressure. in-vacuo – 25-75 °C; 0.05 to 1 mm

5 Unless otherwise indicated, "the residue is chromatographed" is defined as silica gel chromatography of residue with moderate nitrogen pressure (flash chromatography) or a medium pressure chromatography systems using a silica gel to crude product ratio of ~10-100.

For HPLC, the conditions listed are for the analytical trace only. For  
10 Preparative HPLC, the eluent is similar to analytical HPLC eluent.

Thin layer chromatography is performed with silica gel plates with UV and/or appropriate staining solution.

NMR spectra are obtained with either 300 or 400 MHz spectrometer.

NMR data is listed to denote spectrum is consistent with assigned structure.

15 "NMR" notation without data denotes spectrum is consistent with assigned structure.

HRMS – high resolution mass spectrum

ES-MS – electrospray mass spectrum

20 Abbreviations:

Aq - aqueous

d – day

eq - equivalent

h – hour

25 m – minute

satd – saturated

disp – dispersion

quant – quantitative

rt for retention time (both small caps to minimize confusion with RT)

30 RT – room temperature

-63-

## Chemical Definitions:

|    |   |
|----|---|
|    | BF <sub>3</sub> -OEt <sub>2</sub> – boron trifluoride etherate        |
|    | BnBr – benzyl bromide   |
|    | CH <sub>2</sub> Cl <sub>2</sub> -dichloromethane                      |
| 5  | CH <sub>3</sub> CN – acetonitrile                                     |
|    | CO-carbon monoxide  |
|    | CsF – cesium fluoride   |
|    | DMAP - 4-(dimethylamino)pyridine                                      |
|    | DMF – N,N-dimethylformamide   |
| 10 | DMSO – dimethylsulfoxide  |
|    | DPPB – 1,4-bis(diphenylphosphino)butane                               |
|    | DPPF – dichloro[1,1'-bis(diphenylphosphino)ferrocene]                 |
|    | EDCI – 3-Ethyl-1-[3-(dimethylamino)propyl]carbodiimide hydrochloride  |
|    | Et <sub>3</sub> N – triethylamine                                     |
| 15 | EtMgBr- ethyl magnesium bromide                                       |
|    | EtOAc – ethyl acetate   |
|    | EtOH – ethanol  |
|    | H <sub>2</sub> – hydrogen pressure                                    |
|    | H <sub>2</sub> NCH <sub>2</sub> CO <sub>2</sub> Me – methyl glycinate |
| 20 | Hept – heptane  |
|    | Hex - hexanes   |
|    | HN(OMe)Me – N-methyl-O-methyl hydroxylamine                           |
|    | HNMe <sub>2</sub> – dimethyl amine                                    |
|    | HATU – O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium        |
| 25 | hexafluorophosphate   |
|    | HOAT – 7-aza-1-hydroxybenzotriazole                                   |
|    | HOBT – 1-hydroxybenzotriazole   |
|    | K <sub>2</sub> CO <sub>3</sub> – potassium carbonate                  |
|    | KI – potassium iodine   |
| 30 | KOH – potassium hydroxide   |
|    | LAH – lithium aluminum hydride  |
|    | LiHMDS – lithium hexamethyldisilazide                                 |



-64-

|    |   |
|----|---|
|    | Lindlar catalyst – Pd-CaCO <sub>3</sub> -PbO                        |
|    | LiOH – lithium hydroxide  |
|    | mCPBA – meta-chloroperbenzoic acid                                  |
|    | MeI – methyl iodide   |
| 5  | MeOH – methanol   |
|    | NaBH <sub>4</sub> – sodium borohydride                              |
|    | MgSO <sub>4</sub> – magnesium sulfate                               |
|    | NaH – sodium hydride  |
|    | NaHCO <sub>3</sub> – sodium bicarbonate                             |
| 10 | NaI – sodium iodide   |
|    | Na <sub>2</sub> SO <sub>4</sub> – sodium sulfate                    |
|    | NH <sub>4</sub> Cl – ammonium chloride                              |
|    | NMO – 4-methylmorpholine N-oxide                                    |
|    | NMP – N-methylpyrrolidin-2-one                                      |
| 15 | Na-S-R <sub>3</sub> – sodium alkylmercaptide                        |
|    | PBr <sub>3</sub> – phosphorus tribromide                            |
|    | Pd(DPPF) – palladium dichloro[1,1'-bis(diphenylphosphino)ferrocene] |
|    | Pd(OAc) <sub>2</sub> – palladium (II) acetate                       |
|    | Pd(TPP) <sub>4</sub> – palladium tetrakis(triphenylphosphine)       |
| 20 | Pd-C – palladium on carbon  |
|    | Pd-C/H <sub>2</sub> – palladium on carbon with hydrogen pressure    |
|    | pTSA – para-toluenesulfonic acid                                    |
|    | Pyr – pyridine  |
|    | Red-Al – sodium bis(2-methoxyethoxy)aluminum hydride                |
| 25 | R <sub>2</sub> MgBr – alkyl magnesium bromide                       |
|    | R <sub>3</sub> MgBr – alkyl magnesium bromide                       |
|    | R <sub>5</sub> MgBr – alkyl magnesium bromide                       |
|    | R <sub>3</sub> S(O) <sub>2</sub> Cl – alkylsulfonyl chloride        |
|    | R <sub>2</sub> S(O) <sub>2</sub> NH <sub>2</sub> – alkylsulfonamide |
| 30 | TBSCl – <i>tert</i> -butyldimethylsilyl chloride                    |
|    | tBuC(O)CH <sub>2</sub> Br – 1-bromopinacolone                       |
|    | Tf <sub>2</sub> O – triflic anhydride                               |

-65-

TFA – trifluoroacetic acid

THF – tetrahydrofuran

Ti(OiPr)<sub>4</sub> – titanium tetraisopropoxide


TMS-acetylene – trimethylsilyl acetylene

5. TPAP – tetrapropylammonium perruthenate


Zn(OTf)<sub>2</sub> – zinc trifluoromethane sulfonate

## Example 1

Preparation of 2-(4-{1-Ethyl-1-[4-(1-hydroxy-cyclopentylethynyl)-3-methyl-phenyl]-propyl}-2-methyl-benzoylamino)-2-methyl-propionic acid.



A. [E, Z]-3-(3-Methyl-4-hydroxyphenyl)-3-pentene




A mixture of 3-(3-methyl-4-hydroxyphenyl)-3-pentanol (20.0 g, 102.9 mmol),  
 15 pTSA monohydrate (50 mg, 0.26 mmol catalytic amount), cyclohexane (150 mL), and  
 toluene (250 mL) is refluxed on a steam bath for 3 h. During the reflux time, water is  
 separated by means of a Dean-Stark trap. Analysis of the reaction mixture by TLC  
 (SiO<sub>2</sub>; CHCl<sub>3</sub>) shows a spot to spot transformation of the starting material into a less  
 polar material. The cyclohexane-toluene solution is cooled to RT, washed with satd  
 20 sodium carbonate solution (25 mL), MgSO<sub>4</sub> dried, and concentrated to give the title  
 compounds as a [E:Z] isomeric mixture of [4:1] (17.36 g, 96%).

H-NMR (300 MHz, DMSO-d<sub>6</sub>): δ 6.85-7.10 (3H, m), 5.56 (0.8H, q, J = 6.8 Hz), 5.39,  
 0.2H, q, J = 6.8 Hz), 2.40( (1.6H, q, J = 7.6 Hz), 2.25 (0.4H, q, J = 7.6 Hz), 2.15  
 (0.6H, s), 2.05 (2.4H, s), 1.70 (2.4H, d, J = 7.6 Hz), 1.51 (0.6H, d, J = 7.6 Hz), 0.90  
 25 (2.4H, t, J = 7.6 Hz), 0.85 (0.4H, t, J = 7.6 Hz).

-66-

ES GC MS  $m/z$  176.1,  $[M+]$ ; Calc.  $C_{12}H_{16}O$   $m/z$  176.1.

B. [E,Z]-3-[3-Methyl-4-(trifluoromethylsulfonyloxy)phenyl]-3-pentene.




5 A solution of [E,Z]-3-(3-methyl-4-hydroxyphenyl)-3-pentene (17.0 g, 96.4 mmol) and triethylamine (10.3 g, 101 mmol) in  $CH_2Cl_2$  (400 mL) under a  $N_2$  atmosphere is cooled to  $-35^\circ C$  in a dry ice bath. Triflic anhydride (28.6 g, 101 mmol) is added slowly by syringe so as to maintain the temperature below  $-30^\circ C$ . The resulting pale yellow solution is stirred for 3 h as it is allowed to warm to RT. The reaction mixture is poured  
10 over 300 ml pH 7.0 buffer and 300 ml ice. The organic layer is separated and the  $CH_2Cl_2$  layer is washed with additional (4x 150 mL) buffer and dried over anhydrous  $MgSO_4$ . Removal of the solvent under reduced pressure gives the crude product as a yellow oil (approx 30 g). The product is purified by chromatography over silica using elution with 5%  $CHCl_3$  in hexane. The desired product is obtained as a clear, colorless liquid as a  
15 mixture of [E:Z] isomers ratio of 4:1, respectively (29.4 g, 99%).

H-NMR (300 MHz,  $DMSO-d_6$ ):  $\delta$  7.01-7.28 (3H, m), 5.76 (0.8H, q,  $J = 6.8$  Hz), 5.57, 0.2H, q,  $J = 6.8$  Hz), 2.52 (1.6H, q,  $J = 7.6$  Hz), 2.25 (0.4H, q,  $J = 7.6$  Hz), 2.40 (0.6H, s), 2.39 (2.4H, s), 1.81 (2.4H, d,  $J = 7.6$  Hz), 1.55 (0.6H, d,  $J = 7.6$  Hz), 0.99 (2.4H, t,  $J = 7.6$  Hz), 0.97 (0.4H, t,  $J = 7.6$  Hz).

20 ES GC MS  $m/z$  308.1,  $[M+]$ ; Calc. for  $C_{13}H_{15}F_3O_3S$   $m/z$  308.1.

C. [E,Z]-3-(3-Methyl-4-carbomethoxyphenyl)-3-pentene.



A mixture of [E,Z]-3-[3-methyl-4-(trifluoromethylsulfonyloxy)phenyl]-3-pentene  
25 (25.0 g, 81 mmol), MeOH, dppb, DMSO,  $Et_3N$ , and  $Pd(OAc)_2$  is carbonylated by reaction at  $80^\circ C$  under an atmosphere of CO (100 psi). for a total of 6 h. The cooled


-67-

reaction mixture is filtered through filter-cel with the aid of additional MeOH . The filtrate is concentrated under reduced pressure to remove most of the DMSO and the residue is distributed between EtOAc and brine. The organic layer is washed 5 times with brine , dried over MgSO<sub>4</sub>, and concentrated to an oil. The oil is purified by

5 chromatography over silica gel using a 10% to 50% chloroform in hexane gradient . Concentration of appropriate fractions provides the desired product as a clear, nearly colorless liquid as a mixture of [E:Z] isomers ratio of 4:1, respectively (15.2 g, 86%). TLC (CHCl<sub>3</sub>): R<sub>f</sub> ~ 0.7

H-NMR (300 MHz, CDCl<sub>3</sub>): δ 7.03-7.93 (3H, m), 5.83 (0.75H, q, J = 6.9 Hz), 5.57  
10 (0.25H, q, J = 6.9 Hz), 3.91 (0.6H, s), 3.90 (2.4H, s), 2.54( 1.5H, q, J = 7.6 Hz), 2.35 (0.5H, q, J = 7.6 Hz), 1.83 (2.25H, d, J = 7.6 Hz), 1.57 (0.75H, d, J = 7.6 Hz), 1.00 (2.25H, t, J = 7.6 Hz ), 0.97 (0.75H, t, J = 7.6 Hz).

D. 3'-[3-Methyl-4-(hydroxy)phenyl]-3'-(3-methyl-4-carbomethoxy-  
15 phenyl)pentane.



A 25 °C solution of [E,Z]-3-[3-methyl-4-(carbomethoxy)phenyl]-3-pentene  
(5.00g, 22.9 mmol) in o-cresol (9.90 g, 91.6 mmol) is treated with BF<sub>3</sub>-etherate (5.6 g,  
39.5 mmol) and the reaction mixture is stirred for 16 h at ambient temperature. Analysis  
20 by TLC (SiO<sub>2</sub>; CHCl<sub>3</sub>) shows loss of the starting material and formation of a major  
product spot at slightly lower R<sub>f</sub> than o-cresol. The reaction mixture is taken up in 200  
mL ice water and 200 mL ether and carefully basified with excess solid NaHCO<sub>3</sub>. The  
ether layer is separated, washed with brine, dried over anhydrous MgSO<sub>4</sub> and  
concentrated to an oil. The oil is vacuum distilled with 3 x 15 mL ethylene glycol to  
25 remove the remaining o-cresol and the resulting pot residue is cooled and redistributed  
into ether and water and treated with NaHCO<sub>3</sub>, as above. The ether layer is separated,  
washed with water, dried over MgSO<sub>4</sub> and concentrated to provide the crude product as a  
yellow oil (7.3 g). The oil is purified by chromatography over silica gel using CHCl<sub>3</sub> to  
provide the title compound as a pale yellow oil (5.42 g, 72%).

-68-


H-NMR (300 MHz, CDCl<sub>3</sub>): 6.66 to 7.83 (6H, m) 4.54 (1H, s), 3.89 (3H, s), 2.61 (3H, s), 2.21 (3H, s), 2.08 (4H, q, J = 7.3 Hz), 0.62 (6H, t, J = 7.3 Hz).

ES (-) MS m/z 325.2 [M-H]

IR (CHCl<sub>3</sub>): 1716 cm<sup>-1</sup>

5


E. 4-[1-Ethyl-1-(3-methyl-4-trifluoromethanesulfonyloxy-phenyl)-propyl]-2-methyl-benzoic acid methyl ester



4-[1-Ethyl-1-(4-hydroxy-3-methyl-phenyl)-propyl]-2-methyl-benzoic acid methyl ester (10.0 g, 30.63 mmol) in pyridine (30 ml) is added dropwise trifluoromethane sulfonic anhydride (10.37 g, 36.76 mmol) and warmed to room temperature overnight. The reaction is quenched by pouring into ice/water mix (200 ml) and extracted with Et<sub>2</sub>O (3 x 200 ml). The combined organic layers are washed with water (100 ml), 1N HCl (100 ml), water (100 ml), brine (100 ml), dried (MgSO<sub>4</sub>), concentrated and chromatographed (330 g SiO<sub>2</sub>, 5% EtOAc/Hexanes) to yield the title compound (12.46 g, 90%) as a pale yellow oil.

<sup>1</sup>NMR (400MHz, CDCl<sub>3</sub>) δ ppm: 7.85 (d, 1H, J=8.4 Hz), 7.15 (d, 1H, J=8.4 Hz), 7.10-7.02 (m, 4H), 7.10-7.02 (m, 4H), 3.91 (s, 3H), 2.60 (s, 3H), 2.35 (s, 3H), 2.13 (q, 4H, J=7.3 Hz), 0.64 (t, 6H, J=7.3 Hz).

F. 4-[1-Ethyl-1-(3-methyl-4-trimethylsilanylethynyl-phenyl)-propyl]-2-methyl-benzoic acid methyl ester



4-[1-Ethyl-1-(3-methyl-4-trifluoromethanesulfonyloxy-phenyl)-propyl]-2-methyl-benzoic acid methyl ester (11.20 g, 24.43 mmol) in DMF (100 ml) is added trimethylsilylacetylene (6.90 ml, 48.85 mmol), dichlorobis(triphenylphosphine)


25

-69-

palladium (1.71 g, 2.44 mmol) and triethylamine (23.84 ml, 171 mmol) and heated to 80 °C overnight. The reaction mixture is concentrated and filtered through a pad of silica gel eluted with 10% EtOAc/Hexanes. The eluent is concentrated and chromatographed (330 g SiO<sub>2</sub>, 2% EtOAc/Hexanes) to yield the title compound (8.07 g, 81%) as an orange oil.

- 5 <sup>1</sup>NMR (400MHz, CDCl<sub>3</sub>) δ ppm: 7.82 (d, 1H, J=7.9 Hz), 7.35 (d, 1H, J=8.4 Hz), 7.08-6.97 (m, 3H), 6.94 (d, 1H, J=8.4 Hz), 3.90 (s, 3H), 2.57 (s, 3H), 2.41 (s, 3H), 2.11 (q, 5H, J=7.7 Hz), 0.62 (t, 6H, J=7.3 Hz), 0.28 (s, 9H).


- 10 G. 4-[1-Ethyl-1-(4-ethynyl-3-methyl-phenyl)-propyl]-2-methyl-benzoic acid methyl ester



- To an ambient temperature suspension of 4-[1-Ethyl-1-(3-methyl-4-trimethylsilanylethynyl-phenyl)-propyl]-2-methyl-benzoic acid methyl ester (6.8 g, 16.72 mmol) in acetonitrile/water (180/30 ml) is added cesium fluoride (25.4 g, 167.2 mmol) and stirred for 3d. The reaction mixture is concentrated and partitioned between EtOAc (500 ml) and 1N HCl (200 ml). The aqueous phase is extracted with a second portion of EtOAc (200 ml) and the combined organic phases are washed with brine (200 ml), dried (MgSO<sub>4</sub>), concentrated and chromatographed (330 g SiO<sub>2</sub>, 1% EtOAc/Hexanes) to yield the title compound (5.52 g, 99%).
- 20 <sup>1</sup>NMR (400MHz, CDCl<sub>3</sub>) δ ppm: 7.80 (d, 1H, J=8.4 Hz), 7.34 (d, 1H, J=8.4 Hz), 7.04-6.99 (m, 2H), 6.97 (s, 1H), 6.92 (d, 1H, J=7.9 Hz), 3.86 (s, 3H), 3.23 (s, 1H), 2.55 (s, 3H), 2.39 (s, 3H), 2.09 (q, 4H, J=7.2 Hz), 0.60 (t, 6H, J=7.3 Hz).

- 25 H. 4-{1-Ethyl-1-[4-(1-hydroxy-cyclopentylethynyl)-3-methyl-phenyl]-propyl}-2-methyl-benzoic acid methyl ester


-70-



To a  $-78^{\circ}\text{C}$  solution of 4-[1-Ethyl-1-(4-ethynyl-3-methyl-phenyl)-propyl]-2-methyl-benzoic acid methyl ester (2.50 g, 7.47 mmol) in THF (80 ml) is added lithium hexamethyldisilazide (8.22 ml, 8.22 mmol, 1.0 M in THF) and stirred at  $-78^{\circ}\text{C}$  for 30 min. Cyclopentanone (730  $\mu\text{L}$ , 8.22 mmol) is added and the reaction mixture warmed to room temperature. The reaction is quenched with saturated aqueous  $\text{NH}_4\text{Cl}$  (30 ml) and concentrated. The residue is partitioned between EtOAc (300 ml) and 1.0N HCl (50 ml). The aqueous layer is extracted with EtOAc (100 ml). The combined organic layer are washed with water (50 ml), brine (50 ml), dried ( $\text{MgSO}_4$ ), concentrated and chromatographed (120 g  $\text{SiO}_2$ , 5% EtOAc/Hexanes) to yield the title compound (2.45 g, 79%).

$^1\text{NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  ppm: 7.83 (d, 1H,  $J=8.8$  Hz), 7.32 (dd, 1H,  $J=7.9, 2.2$  Hz), 7.08-6.98 (m, 3H), 6.95 (d, 1H,  $J=8.4$  Hz), 3.91 (s, 3H), 2.58 (s, 3H), 2.41 (s, 3H), 2.12 (q, 4H,  $J=7.2$  Hz), 1.87-1.74 (m, 4H), 1.15 (t, 6H,  $J=7.4$  Hz), 0.64 (t, 6H,  $J=7.3$  Hz).

I. 4-{1-Ethyl-1-[4-(1-hydroxy-cyclopentylethynyl)-3-methyl-phenyl]-propyl}-2-methyl-benzoic acid.



To an ambient temperature solution of 4-{1-Ethyl-1-[4-(1-hydroxy-cyclopentylethynyl)-3-methyl-phenyl]-propyl}-2-methyl-benzoic acid methyl ester (2.40 g, 5.73 mmol) in dioxane/water (30/30 ml) is added lithium hydroxide (412 mg, 17.20 mmol) and heated to  $60^{\circ}\text{C}$  for 1h. The reaction is concentrated and partitioned between  $\text{Et}_2\text{O}$ /EtOAc (150/150 ml) and 1N HCl (50 ml). The aqueous phase is extracted with a


-71-

second portion of EtOAc (100 ml) and the combined organic phases are washed with water (50 ml), brine (50 ml), dried ( $\text{MgSO}_4$ ) and concentrated to yield the title compound (1.57 g, 68%).

$^1\text{NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  ppm: 7.93 (d, 1H,  $J=8.4$  Hz), 7.28 (d, 1H,  $J=8.4$  Hz), 7.05 (d, 1H,  $J=8.4$  Hz), 7.03 (s, 1H), 6.96 (s, 1H), 6.92 (dd, 1H,  $J=8.1, 2.0$  Hz), 2.58 (s, 3H), 2.36 (s, 3H), 2.15-1.97 (m, 8H), 1.94-1.72 (m, 4H), 0.60 (t, 6H,  $J=7.3$  Hz).

High Res. EI-MS: 403.2270; calc. for  $\text{C}_{27}\text{H}_{32}\text{O}_3\text{-H}$ : 403.2273.

J. 2-(4-{1-Ethyl-1-[4-(1-hydroxy-cyclopentylethynyl)-3-methyl-phenyl]-propyl}-2-methyl-benzoylamino)-2-methyl-propionic acid methyl ester.



4-{1-Ethyl-1-[4-(1-hydroxy-cyclopentylethynyl)-3-methyl-phenyl]-propyl}-2-methyl-benzoic acid (389 mg, 0.962 mmol),  $\alpha$ -aminoisobutyric acid methyl ester hydrochloride (221 mg, 1.44 mmol), 1-(3-dimethyl-aminopropyl)-3-ethylcarbodiimide hydrochloride (276 mg, 1.44 mmol), 1-hydroxybenzotriazole hydrate (195 mg, 1.44 mmol) and triethylamine (536  $\mu\text{L}$ , 3.85 mmol) are stirred overnight. The reaction is concentrated and partitioned between EtOAc (150 ml) and 1N HCl (50 ml). The aqueous layer is extracted with EtOAc (100 ml). The combined organic layers are washed with water (50 ml),  $\text{NaHCO}_3$  (50 ml), brine (50 ml), dried (anhydrous  $\text{MgSO}_4$ ), concentrated and chromatographed (12 g  $\text{SiO}_2$ , 15% EtOAc/Hexanes) to yield the title compound (398 mg, 82%).


$^1\text{NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  ppm: 7.29-7.24 (m, 2H), 6.99-6.88 (m, 4H), 6.27 (s, 1H), 3.77 (s, 3H), 2.39 (s, 3H), 2.36 (s, 3H), 2.12-1.98 (m, 8H), 1.93-1.73 (m, 4H), 1.65 (s, 6H), 0.59 (t, 6H,  $J=7.3$  Hz).

High Res. EI-MS: 504.3133; calc. for  $\text{C}_{32}\text{H}_{41}\text{NO}_4\text{-H}$ : 504.3114.

K. 2-(4-{1-Ethyl-1-[4-(1-hydroxy-cyclopentylethynyl)-3-methyl-phenyl]-propyl}-2-methyl-benzoylamino)-2-methyl-propionic acid.



-72-




2-(4-{1-Ethyl-1-[4-(1-hydroxy-cyclopentylethynyl)-3-methyl-phenyl]-propyl}-2-methyl-benzoylamino)-2-methyl-propionic acid methyl ester (360 mg, 0.715 mmol) and lithium hydroxide (51 mg, 2.15 mmol) are reacted and purified in a procedure analogous to Example 11 to give the title compound (353 mg, quant.).

$^1\text{NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  ppm: 7.27 (d, 2H,  $J=8.4$  Hz), 7.01-6.87 (m, 4H), 6.22 (s, 1H), 2.40 (s, 3H), 2.36 (s, 3H), 2.12-1.98 (m, 8H), 1.93-1.73 (m, 4H), 1.68 (s, 6H), 0.59 (t, 6H,  $J=7.0$  Hz).


High Res. EI-MS: 490.2957; calc. for  $\text{C}_{31}\text{H}_{39}\text{NO}_4+\text{H}$ : 490.2975.

#### Example 2.

Preparation of 2-(4-{1-Ethyl-1-[4-(3-ethyl-3-hydroxy-pent-1-ynyl)-3-methyl-phenyl]-propyl}-2-methyl-benzoylamino)-2-methyl-propionic acid.



A. 4-{1-Ethyl-1-[4-(3-ethyl-3-hydroxy-pent-1-ynyl)-3-methyl-phenyl]-propyl}-2-methyl-benzoic acid methyl ester.




4-[1-Ethyl-1-(4-ethynyl-3-methyl-phenyl)-propyl]-2-methyl-benzoic acid methyl ester (2.50 g, 7.47 mmol), lithium hexamethyldisilazide (8.22 ml, 8.22 mmol, 1.0 M in

-73-

THF) and 3-pentanone (870  $\mu$ L, 8.22 mmol) are reacted and purified analogous to Example 1H to give the title compound (2.22 g, 71%).

$^1$ NMR (400MHz,  $\text{CDCl}_3$ )  $\delta$  ppm: 7.79 (d, 1H,  $J=7.9$  Hz), 7.27 (d, 1H,  $J=8.6$  Hz), 7.04-6.98 (m, 2H), 6.95 (s, 1H), 6.91 (dd, 1H,  $J=8.0, 1.9$  Hz), 3.86 (s, 3H), 2.54 (s, 3H), 2.36 (s, 3H), 2.13-1.98 (m, 8H), 1.94-1.72 (m, 5H), 0.59 (t, 6H,  $J=7.3$  Hz).

B. 4-{1-Ethyl-1-[4-(3-ethyl-3-hydroxy-pent-1-ynyl)-3-methyl-phenyl]-propyl}-2-methyl-benzoic acid.




10 4-{1-Ethyl-1-[4-(3-ethyl-3-hydroxy-pent-1-ynyl)-3-methyl-phenyl]-propyl}-2-methyl-benzoic acid methyl ester (2.20 g, 5.23 mmol) and lithium hydroxide (376 mg, 15.69 mmol) are reacted and purified analogous to Example 1I to give the title compound (1.77 g, 83%).

$^1$ NMR (400MHz,  $\text{CDCl}_3$ )  $\delta$  ppm: 7.93 (d, 1H,  $J=7.9$  Hz), 7.29 (d, 1H,  $J=8.4$  Hz), 7.08-7.01 (m, 2H), 6.97 (s, 1H), 6.92 (d, 1H,  $J=7.5$  Hz), 2.59 (s, 3H), 2.38 (s, 3H), 2.10 (q, 4H,  $J=7.3$  Hz), 1.84-1.69 (m, 4H), 1.11 (t, 6H,  $J=7.2$  Hz), 0.61 (t, 6H,  $J=7.3$  Hz).

High Res. EI-MS: 405.2444; calc. for  $\text{C}_{27}\text{H}_{34}\text{O}_3\text{-H}$ : 405.2430.

20 C. 2-(4-{1-Ethyl-1-[4-(3-ethyl-3-hydroxy-pent-1-ynyl)-3-methyl-phenyl]-propyl}-2-methyl-benzoylamino)-2-methyl-propionic acid methyl ester.



4-{1-Ethyl-1-[4-(3-ethyl-3-hydroxy-pent-1-ynyl)-3-methyl-phenyl]-propyl}-2-methyl-benzoic acid (437 mg, 1.07 mmol),  $\alpha$ -aminoisobutyric acid methyl ester


-74-

hydrochloride (247 mg, 1.61 mmol), 1-(3-dimethyl-aminopropyl)-3-ethylcarbodiimide hydrochloride (309 mg, 1.61 mmol), 1-hydroxybenzotriazole hydrate (218 mg, 1.61 mmol) and triethylamine (597  $\mu$ L, 4.28 mmol) are reacted and purified analogous to Example 1J to give the title compound (419 mg, 77%).

- 5  $^1\text{NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  ppm: 7.29-7.25 (m, 2H), 7.00-6.89 (m, 4H), 6.27 (s, 1H), 3.77 (s, 3H), 2.39 (s, 3H), 2.37 (s, 3H), 2.06 (q, 4H,  $J=7.3$  Hz), 1.83-1.70 (m, 4H), 1.65 (s, 6H), 1.11 (t, 6H,  $J=7.5$  Hz), 0.59 (t, 6H,  $J=7.3$  Hz).

High Res. EI-MS: 506.3271; calc. for  $\text{C}_{32}\text{H}_{43}\text{NO}_4+\text{H}$ : 506.3270.

- 10 D. 2-(4-{1-Ethyl-1-[4-(3-ethyl-3-hydroxy-pent-1-ynyl)-3-methyl-phenyl]-propyl}-2-methyl-benzoylamino)-2-methyl-propionic acid.



- 2-(4-{1-Ethyl-1-[4-(3-ethyl-3-hydroxy-pent-1-ynyl)-3-methyl-phenyl]-propyl}-2-methyl-benzoylamino)-2-methyl-propionic acid methyl ester (397 mg, 0.785 mmol) and  
 15 lithium hydroxide (56 mg, 2.36 mmol) are reacted and purified analogous to Example 1I to give the title compound (384 mg, 99%).


- $^1\text{NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  ppm: 7.30-7.26 (m, 2H), 7.02-6.97 (m, 2H), 6.95 (s, 1H), 6.91 (d, 1H,  $J=8.4$  Hz), 6.21 (s, 1H), 2.40 (s, 3H), 2.37 (s, 3H), 2.07 (q, 4H,  $J=6.6$  Hz), 1.83-1.71 (m, 4H), 1.68 (s, 6H), 1.11 (t, 6H,  $J=7.3$  Hz),  
 20 0.59 (t, 6H,  $J=7.3$  Hz).

High Res. EI-MS: 492.3116; calc. for  $\text{C}_{31}\text{H}_{41}\text{NO}_4+\text{H}$ : 492.3114.


### Example 3

- 25 Preparation of (4-{1-ethyl-1-[4-(2-ethyl-2-hydroxy-butoxy)-3-methyl-phenyl]-propyl}-2-methyl-benzoylamino)-acetic acid.

-75-



A. 3',3'-Bis[4-hydroxy-3-methylphenyl]pentane.




- 5 To a mixture of o-cresol (196 g, 1.81 mol) and 3-pentanone (60 ml, 0.57 mol) is added methanesulfonic acid (45 ml, 0.69 mol) and stirred for 3 days. The reaction is basified to pH 8 with satd  $\text{Na}_2\text{CO}_3$  and extracted with EtOAc. The organic layer is washed with water (6 X 500 ml),  $\text{Na}_2\text{SO}_4$  dried, concentrated, chromatographed (2 kg  $\text{SiO}_2$ , Hex to 80% EtOAc/Hex), and triturated with Hex to give the title compound as a white solid (100 g, 61%).

NMR 400 MHz(DMSO):  $\delta$  0.49 (t,  $J = 7.3$  Hz, 6H), 1.91 (q,  $J = 7.3$  Hz, 4H), 2.02 (s, 6H), 6.61 (d,  $J = 8.3$  Hz, 2H), 6.73 (d,  $J = 8.3$  Hz, 2H), 6.76 (s, 2H), 8.94 (s, 2H).

High Res. EI-MS: 284.1794; calc. for  $\text{C}_{19}\text{H}_{24}\text{O}_2$ : 284.1776

- 15 B. 3'-[4-Benzyloxy-3-methylphenyl]-3'-[4-hydroxy-3-methylphenyl]pentane.




-76-

To a solution of 3,3-bis[4-hydroxy-3-methylphenyl]pentane (10 g, 35.2 mmol) and DMF (180 ml) is added 60% NaH disp (1.4 g, 35.2 mmol). After stirring for 30 m, to the reaction is added benzyl bromide (4.2 ml, 35.2 mmol). The mixture is stirred for 14 h and concentrated in vacuo. The residue is partitioned between Et<sub>2</sub>O/water. The organic layer is washed with 1N HCl, water, brine, Na<sub>2</sub>SO<sub>4</sub> dried, concentrated, and chromatographed (MeCl<sub>2</sub>) to give the title compound as an oil (6.5 g, 49%)

High Res. FAB-MS: 374.2237; calc. for C<sub>26</sub>H<sub>30</sub>O<sub>2</sub>: 374.2246

C. {4-[1-(4-Benzyloxy-3-methyl-phenyl)-1-ethyl-propyl]-2-methyl-phenoxy}-acetic acid methyl ester.




Using a procedure analogous to Example 3B, from 4-[1-(4-benzyloxy-3-methyl-phenyl)-1-ethyl-propyl]-2-methyl-phenol (9.76 g, 25.93 mmol) and methyl bromoacetate (2.7 mL, 28.53 mmol) gives the title compound (10.24 g, 22.96 mmol, 89%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>), δ 0.61 (t, *J* = 7.1 Hz, 6H), 2.04 (q, *J* = 7.1 Hz, 4H), 2.23 (s, 3H), 2.24 (s, 3H), 3.80 (s, 3H), 4.62 (s, 2H), 5.05 (s, 2H), 6.58 (d, *J* = 7.8 Hz, 1H), 6.77 (d, *J* = 8.4 Hz, 1H), 6.90-6.97 (m, 4H), 7.29-7.48 (m, 5H).

ES-MS (*m/z*): calcd for C<sub>29</sub>H<sub>38</sub>NO<sub>4</sub> (M+NH<sub>4</sub>)<sup>+</sup>: 464.6; found: 464.3.

-77-

D. 3-{4-[1-(4-Benzyloxy-3-methyl-phenyl)-1-ethyl-propyl]-2-methyl-phenoxy-methyl}-pentan-3-ol.




5 To a solution of {4-[1-(4-benzyloxy-3-methyl-phenyl)-1-ethyl-propyl]-2-methyl-phenoxy}-acetic acid methyl ester (4.78 g, 10.72 mmol) in THF (40 mL) at 0 °C is added 3.0 M EtMgBr (8.9 mL, 26.8 mmol) in a dropwise fashion. After the addition, it is stirred at 0 °C for 10 m, then refluxed for 3 h. The reaction mixture is poured into ice-H<sub>2</sub>O, acidified with 0.1 M HCl and extracted with EtOAc (100 mL);  
 10 MgSO<sub>4</sub> dried, and concentrated and chromatographed to give the title compound (4.06 g, 8.56 mmol, 80%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>), δ 0.62 (t, *J* = 7.4 Hz, 6H), 0.95 (t, *J* = 7.4 Hz, 6H), 1.68 (q, *J* = 7.4 Hz, 2H), 1.69 (q, *J* = 7.4 Hz, 2H), 2.00-2.09 (m, 4H), 2.19 (s, 3H), 2.23 (s, 3H), 3.81 (s, 2H), 5.05 (s, 2H), 6.70 (d, *J* = 8.4 Hz, 1H), 6.77 (d, *J* = 7.8 Hz, 1H), 6.90-6.99 (m,  
 15 4H), 7.29-7.48 (m, 5H).

ES-MS (*m/z*): calcd for C<sub>32</sub>H<sub>46</sub>NO<sub>3</sub> (M+NH<sub>4</sub>)<sup>+</sup>: 492.7; found: 492.4.

E. Acetic acid 1-{4-[1-(4-benzyloxy-3-methyl-phenyl)-1-ethyl-propyl]-2-methyl-phenoxy-methyl}-1-ethyl-propyl ester.

20




-78-

A reaction mixture containing 3-{4-[1-(4-benzyloxy-3-methyl-phenyl)-1-ethyl-propyl]-2-methyl-phenoxy-methyl}-pentan-3-ol (3.92 g, 8.27 mmol), Et<sub>3</sub>N (11.5 mL, 82.7 mmol) and Ac<sub>2</sub>O (3.9 mL, 41.3 mmol), is stirred at rt for 3 h. It is diluted with EtOAc (100 mL), washed with 0.1 M HCl (2 x 60 mL) and brine (60 mL); MgSO<sub>4</sub> dried and concentrated to give the title compound (2.82 g, 5.47 mmol, 66%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>), δ 0.62 (t, *J* = 7.5 Hz, 6H), 0.91 (t, *J* = 7.5 Hz, 6H), 1.98-2.09 (m, 11H), 2.16 (s, 3H), 2.24 (s, 3H), 4.13 (s, 2H), 5.05 (s, 2H), 6.69 (d, *J* = 8.4 Hz, 1H), 6.77 (d, *J* = 9.2 Hz, 1H), 6.90-6.98 (m, 4H), 7.29-7.47 (m, 5H).

ES-MS (*m/z*): calcd for C<sub>34</sub>H<sub>48</sub>NO<sub>4</sub> (M+NH<sub>4</sub>)<sup>+</sup>: 534.7; found: 534.4.

F. Acetic acid 1-ethyl-1-{4-[1-ethyl-1-(4-hydroxy-3-methyl-phenyl)-propyl]-2-methyl-phenoxy-methyl}-propyl ester.




Acetic acid 1-{4-[1-(4-benzyloxy-3-methyl-phenyl)-1-ethyl-propyl]-2-methyl-phenoxy-methyl}-1-ethyl-propyl ester (0.14 g, 0.27 mmol), EtOH (20 mL), and 10% Pd/C (25 mg) is hydrogenated at atmospheric pressure for 18 h. The reaction is filtered through diatomaceous earth with EtOAc wash. The filtrate is concentrated to give the title compound (0.11 g, 0.25 mmol, 94%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>), δ 0.61 (t, *J* = 7.0 Hz, 6H), 0.91 (t, *J* = 7.5 Hz, 6H), 1.97-2.09 (m, 11H), 2.16 (s, 3H), 2.21 (s, 3H), 4.13 (s, 2H), 6.65 (d, *J* = 8.3 Hz, 1H), 6.69 (d, *J* = 8.3 Hz, 1H), 6.84-6.96 (m, 4H).

ES-MS (*m/z*): calcd for C<sub>27</sub>H<sub>37</sub>O<sub>4</sub> (M-H)<sup>-</sup>: 425.6; found: 425.4.

-79-

D. Acetic acid 1-ethyl-1-{4-[1-ethyl-1-(3-methyl-4-trifluoromethanesulfonyloxy-phenyl)-propyl]-2-methyl-phenoxy-methyl}-propyl ester.




5 Using a procedure analogous to Example 1B, acetic acid 1-ethyl-1-{4-[1-ethyl-1-(4-hydroxy-3-methyl-phenyl)-propyl]-2-methyl-phenoxy-methyl}-propyl ester (2.32 g, 5.43 mmol) gives the title compound (1.94 g, 3.48 mmol, 64%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>), δ 0.61 (t, *J* = 7.5 Hz, 6H), 0.90 (t, *J* = 7.5 Hz, 6H), 1.97-2.10 (m, 11H), 2.15 (s, 3H), 2.56 (s, 3H), 4.13 (s, 2H), 6.69 (d, *J* = 8.3 Hz, 1H), 6.83 (d, *J* = 1.8 Hz, 1H),  
 10 6.92 (dd, *J* = 8.4, 2.2 Hz, 1H), 7.03-7.05 (m, 2H), 7.79 (d, *J* = 7.9 Hz, 1H). ES-MS (*m/z*): calcd for C<sub>28</sub>H<sub>41</sub>F<sub>3</sub>NO<sub>6</sub>S (M+NH<sub>4</sub>)<sup>+</sup>: 576.7; found: 576.3.

E. 4-{1-[4-(2-Acetoxy-2-ethyl-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-2-methyl-benzoic acid methyl ester.

15



Using a procedure analogous to Example 1C, acetic acid 1-ethyl-1-{4-[1-ethyl-1-(3-methyl-4-trifluoromethanesulfonyloxy-phenyl)-propyl]-2-methyl-phenoxy-methyl}-propyl ester (1.94 g, 3.48 mmol) gives the title compound (1.37 g, 2.95 mmol, 85%).


20 <sup>1</sup>H NMR (CDCl<sub>3</sub>), δ 0.61 (t, *J* = 7.5 Hz, 6H), 0.90 (t, *J* = 7.5 Hz, 6H), 1.97-2.11 (m, 11H), 2.15 (s, 3H), 2.56 (s, 3H), 3.87 (s, 3H), 4.13 (s, 2H), 6.69 (d, *J* = 8.8 Hz, 1H), 6.83 (d, *J* =



-80-

1.7 Hz, 1H), 6.92 (dd,  $J = 8.4, 2.2$  Hz, 1H), 7.02-7.07 (m, 2H), 7.79 (d,  $J = 7.9$  Hz, 1H).  
 ES-MS ( $m/z$ ): calcd for  $C_{29}H_{40}O_5$ : 468.6; found: 468.0.

F. 4-{1-Ethyl-1-[4-(2-ethyl-2-hydroxy-butoxy)-3-methyl-phenyl]-propyl}-2-methyl-  
 5 benzoic acid.




Using a procedure analogous to Example 1I, 4-{1-[4-(2-acetoxy-2-ethyl-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-2-methyl-benzoic acid methyl ester (1.38 g, 2.95 mmol) gives the title compound (0.96 g, 2.34 mmol, 79%).

10  $^1H$  NMR ( $CDCl_3$ ),  $\delta$  0.63 (t,  $J = 7.0$  Hz, 6H), 0.95 (t,  $J = 7.0$  Hz, 6H), 1.68 (q,  $J = 7.0$  Hz, 2H), 1.69 (q,  $J = 7.0$  Hz, 2H), 2.10 (q,  $J = 7.0$  Hz, 4H), 2.19 (s, 3H), 2.61 (s, 3H), 3.81 (s, 2H), 6.71 (d,  $J = 8.4$  Hz, 1H), 6.87 (d,  $J = 1.7$  Hz, 1H), 6.94 (dd,  $J = 8.4, 2.3$  Hz, 1H), 7.06-7.10 (m, 2H), 7.94 (d,  $J = 8.8$  Hz, 1H).

ES-MS ( $m/z$ ): calcd for  $C_{26}H_{35}O_4$  ( $M-H$ ): 411.6; found: 411.3.

15

G. (4-{1-Ethyl-1-[4-(2-ethyl-2-hydroxy-butoxy)-3-methyl-phenyl]-propyl}-2-methyl-benzoylamino)-acetic acid methyl ester.



Using a procedure analogous to Example 1J, 4-{1-ethyl-1-[4-(2-ethyl-2-hydroxy-butoxy)-3-methyl-phenyl]-propyl}-2-methyl-benzoic acid (0.64 g, 1.55 mmol) and glycine methyl ester hydrochloride (195 mg, 1.55 mmol) gives the title compound (0.72 g, 1.49 mmol, 96%).


20

-81-

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ),  $\delta$  0.62 (t,  $J = 7.0$  Hz, 6H), 0.95 (t,  $J = 7.0$  Hz, 6H), 1.68 (q,  $J = 7.0$  Hz, 2H), 1.69 (q,  $J = 7.0$  Hz, 2H), 2.07 (q,  $J = 7.0$  Hz, 4H), 2.19 (s, 3H), 2.44 (s, 3H), 3.80 (s, 3H), 3.81 (s, 2H), 4.24 (d,  $J = 5.4$  Hz, 2H), 6.28 (t,  $J = 5.4$  Hz, 1H), 6.70 (d,  $J = 8.3$  Hz, 1H), 6.85 (d,  $J = 2.2$  Hz, 1H), 6.94 (dd,  $J = 8.4, 2.2$  Hz, 1H), 6.99-7.05 (m, 2H), 7.31 (d,  $J = 7.9$  Hz, 1H).

ES-MS ( $m/z$ ): calcd for  $\text{C}_{29}\text{H}_{42}\text{NO}_5$  ( $\text{M}+\text{H}$ ) $^+$ : 484.6; found: 484.4.

H. (4-{1-Ethyl-1-[4-(2-ethyl-2-hydroxy-butoxy)-3-methyl-phenyl]-propyl}-2-methyl-benzoylamino)-acetic acid.




Using a procedure analogous to Example 1I, (4-{1-ethyl-1-[4-(2-ethyl-2-hydroxy-butoxy)-3-methyl-phenyl]-propyl}-2-methyl-benzoylamino)-acetic acid methyl ester (0.72 g, 1.48 mmol) gives the title compound (0.45 g, 0.97 mmol, 657%).

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ),  $\delta$  0.62 (t,  $J = 7.4$  Hz, 6H), 0.95 (q,  $J = 7.4$  Hz, 6H), 1.68 (q,  $J = 7.4$  Hz, 2H), 1.69 (q,  $J = 7.4$  Hz, 2H), 2.06 (q,  $J = 7.4$  Hz, 4H), 2.12 (s, 3H), 2.43 (s, 3H), 3.81 (s, 2H), 4.26 (d,  $J = 5.2$  Hz, 2H), 6.35 (t,  $J = 5.2$  Hz, 1H), 6.70 (d,  $J = 8.4$  Hz, 1H), 6.84 (d,  $J = 1.8$  Hz, 1H), 6.93 (dd,  $J = 8.4, 2.2$  Hz, 1H), 6.99-7.06 (m, 2H), 7.31 (d,  $J = 8.4$  Hz, 1H). ES-MS ( $m/z$ ): calcd for  $\text{C}_{28}\text{H}_{38}\text{NO}_5$  ( $\text{M}-\text{H}$ ) $^-$ : 468.6; found: 468.4.


#### Example 4

Preparation of (4-{1-Ethyl-1-[4-(2-hydroxy-2-isopropyl-3-methyl-butoxy)-3-methyl-phenyl]-propyl}-2-methyl-benzoylamino)-acetic acid.

-82-




A. {4-[1-(4-Benzyloxy-3-methyl-phenyl)-1-ethyl-propyl]-2-methyl-phenoxy}-acetic acid.



Using a procedure analogous to Example 1I, from {4-[1-(4-benzyloxy-3-methyl-phenyl)-1-ethyl-propyl]-2-methyl-phenoxy}-acetic acid methyl ester (1.58 g, 3.54 mmol) gives the title compound (1.47 g, 3.40 mmol, 96%).

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ),  $\delta$  0.61 (t,  $J = 7.5$  Hz, 6H), 2.04 (q,  $J = 7.5$  Hz, 4H), 2.23 (s, 3H), 2.24 (s, 3H), 4.67 (s, 2H), 5.05 (s, 2H), 6.62 (d,  $J = 8.8$  Hz, 1H), 6.77 (d,  $J = 8.3$  Hz, 1H), 6.90-6.98 (m, 4H), 7.29-7.47 (m, 5H). ES-MS ( $m/z$ ): calcd for  $\text{C}_{28}\text{H}_{31}\text{O}_4$  ( $\text{M}-\text{H}$ ): 431.6; found: 431.3.

B. 2-{4-[1-(4-Benzyloxy-3-methyl-phenyl)-1-ethyl-propyl]-2-methyl-phenoxy}-N-methoxy-N-methyl-acetamide.




-83-

Using an analogous procedure to Example 1J, {4-[1-(4-benzyloxy-3-methyl-phenyl)-1-ethyl-propyl]-2-methyl-phenoxy}-acetic acid (1.46 g, 3.38 mmol) and *N,O*-dimethylmethylhydroxyamine hydrochloride (363 mg, 3.72 mmol) give the title compound (1.24 g, 2.61 mmol, 77%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>), δ 0.61 (t, *J* = 7.5 Hz, 6H), 2.03 (q, *J* = 7.5 Hz, 4H), 2.23 (s, 3H), 2.25 (s, 3H), 3.25 (s, 3H), 3.76 (s, 3H), 4.80 (s, 2H), 5.04 (s, 2H), 6.64 (d, *J* = 9.2 Hz, 1H), 6.77 (d, *J* = 8.4 Hz, 1H), 6.90-6.98 (m, 4H), 7.29-7.47 (m, 5H).

ES-MS (*m/z*): calcd for C<sub>30</sub>H<sub>38</sub>NO<sub>4</sub> (M+H)<sup>+</sup>: 476.6; found: 476.4.

C. 1-{4-[1-(4-Benzyloxy-3-methyl-phenyl)-1-ethyl-propyl]-2-methyl-phenoxy}-3-methyl-butan-2-one.



To a 0 °C solution of 2-{4-[1-(4-benzyloxy-3-methyl-phenyl)-1-ethyl-propyl]-2-methyl-phenoxy}-*N*-methoxy-*N*-methyl-acetamide (2.36 g, 4.98 mmol) in THF (50 mL) is added 2.0 M *i*PrMgCl (3.0 mL, 5.98 mmol). The reaction is stirred at 0 °C for 30 m, then at RT for 1 h. It is quenched with satd NH<sub>4</sub>Cl (20 mL), diluted with H<sub>2</sub>O (50 mL),; extracted with EtOAc (2 x 100 mL),; washed with 0.1 M HCl (50 mL), H<sub>2</sub>O (50 mL); MgSO<sub>4</sub> dried, concentrated and purified to give the title compound (1.01 g, 2.21 mmol, 44%).


<sup>1</sup>H NMR (CDCl<sub>3</sub>), δ 0.61 (t, *J* = 7.5 Hz, 6H), 1.17 (d, *J* = 7.0 Hz, 6H), 2.03 (q, *J* = 7.5 Hz, 4H), 2.23 (s, 3H), 2.25 (s, 3H), 3.07 (quintet, *J* = 7.0 Hz, 1H), 4.58 (s, 2H), 5.04 (s, 2H), 6.55 (d, *J* = 8.8 Hz, 1H), 6.77 (d, *J* = 8.4 Hz, 1H), 6.90-6.97 (m, 4H), 7.29-7.47 (m, 5H).

ES-MS (*m/z*): calcd for C<sub>31</sub>H<sub>42</sub>NO<sub>3</sub> (M+NH<sub>4</sub>)<sup>+</sup>: 476.7; found: 476.4.

25


D. 3-{4-[1-(4-Benzyloxy-3-methyl-phenyl)-1-ethyl-propyl]-2-methyl-phenoxy-methyl}-2,4-dimethyl-pentan-3-ol.

-84-



Anhydrous  $\text{CeCl}_3$  (0.82 g) is suspended in THF (30 mL) with stirring overnight. It is cooled to 0 °C, treated with 2.0 M  $\text{iPrMgCl}$  (1.3 mL) and stirred at 0 °C for 1h. A solution of 1-{4-[1-(4-benzyloxy-3-methyl-phenyl)-1-ethyl-propyl]-2-methyl-phenoxy}-3-methyl-butan-2-one (1.01 g, 2.21 mmol) in THF (10 mL) is added. The mixture is stirred at 0 °C for 1 h, before it is quenched with 0.1 M HCl (20 mL). It is extracted with EtOAc (200 mL), washed with 0.1 M HCl (3 x 50 mL);  $\text{MgSO}_4$  dried and concentrated to give the title compound (1.10 g, 2.19 mmol, 99%).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ),  $\delta$  0.62 (t,  $J = 7.0$  Hz, 6H), 1.01 (d,  $J = 7.1$  Hz, 6H), 1.04 (d,  $J = 7.1$  Hz, 6H), 1.57 (b, 2H), 2.04 (q,  $J = 7.0$  Hz, 4H), 2.17 (s, 3H), 2.24 (s, 3H), 3.88 (s, 2H), 5.04 (s, 2H), 6.70 (d,  $J = 8.3$  Hz, 1H), 6.77 (d,  $J = 9.2$  Hz, 1H), 6.90 (d,  $J = 2.2$  Hz, 1H), 6.93-6.98 (m, 3H), 7.29-7.47 (m, 5H). ES-MS ( $m/z$ ): calcd for  $\text{C}_{34}\text{H}_{50}\text{NO}_3$  ( $\text{M} + \text{NH}_4$ ) $^+$ : 520.7; found: 520.63.


15 E. 4-{1-Ethyl-1-[4-(2-hydroxy-2-isopropyl-3-methyl-butoxy)-3-methyl-phenyl]-propyl}-2-methyl-phenol



-85-


Using a procedure analogous to Example 3F, 3-{4-[1-(4-benzyloxy-3-methyl-phenyl)-1-ethyl-propyl]-2-methyl-phenoxy-methyl}-2,4-dimethyl-pentan-3-ol (5.06 g, 10.07 mmol) gives the title compound (3.97 g, 9.64 mmol, 96%).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ),  $\delta$  0.62 (t,  $J = 7.4$  Hz, 6H), 1.00-1.06 (m, 12H), 2.03 (q,  $J = 7.2$  Hz, 4H), 2.11-2.20 (m, 2H),  
 5 2.17 (s, 3H), 2.21 (s, 3H), 3.89 (s, 2H), 6.65 (d,  $J = 8.5$  Hz, 1H), 6.70 (d,  $J = 8.3$  Hz, 1H), 6.87 (dd,  $J = 8.3, 2.2$  Hz, 1H), 6.90 (dd,  $J = 10.0, 2.2$  Hz, 1H), 6.89-6.93 (m, 2H), 6.96 (dd,  $J = 8.5, 2.2$  Hz, 1H).

F. Trifluoro-methanesulfonic acid 4-{1-ethyl-1-[4-(2-hydroxy-2-isopropyl-3-methyl-butoxy)-3-methyl-phenyl]-propyl}-2-methyl-phenyl ester.  
 10



Using a procedure analogous to Example 1B, 4-{1-ethyl-1-[4-(2-hydroxy-2-isopropyl-3-methyl-butoxy)-3-methyl-phenyl]-propyl}-2-methyl-phenol (3.79 g, 9.21 mmol) gives the title compound (1.90 g, 3.49 mmol, 38%).  
 15  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ),  $\delta$  0.62 (t,  $J = 7.2$  Hz, 6H), 1.00-1.06 (m, 12H), 2.05 (q,  $J = 7.4$  Hz, 4H), 2.11-2.20 (m, 2H), 2.18 (s, 3H), 2.33 (s, 3H), 3.89 (s, 2H), 6.62 (d,  $J = 8.4$  Hz, 1H), 6.85 (d,  $J = 2.3$  Hz, 1H), 6.93 (dd,  $J = 8.6, 2.3$  Hz, 1H), 7.04 (dd,  $J = 8.6, 2.1$  Hz, 1H), 7.09 (s, 1H), 7.10 (d,  $J = 6.4$  Hz, 1H).


G. 4-{1-Ethyl-1-[4-(2-hydroxy-2-isopropyl-3-methyl-butoxy)-3-methyl-phenyl]-propyl}-2-methyl-benzoic acid methyl ester.  
 20



-86-


Using a procedure analogous to Example 1C, from trifluoro-methanesulfonic acid 4-{1-ethyl-1-[4-(2-hydroxy-2-isopropyl-3-methyl-butoxy)-3-methyl-phenyl]-propyl}-2-methyl-phenyl ester (3.80 g, 6.98 mmol) to give the title compound (2.75 g, 6.05 mmol, 87%).

- 5  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ),  $\delta$  0.61 (t,  $J = 7.2$  Hz, 6H), 0.98-1.04 (m, 12H), 2.07 (q,  $J = 7.2$  Hz, 4H), 2.10-2.18 (m, 2H), 2.15 (s, 3H), 2.56 (s, 3H), 3.86 (s, 3H), 3.88 (s, 2H), 6.70 (d,  $J = 8.7$  Hz, 1H), 6.84 (d,  $J = 1.6$  Hz, 1H), 6.93 (dd,  $J = 8.5, 2.5$  Hz, 1H), 7.04 (dd,  $J = 8.5, 1.6$  Hz, 1H), 7.06 (s, 1H), 7.79 (d,  $J = 7.8$  Hz, 1H).
- 10 H. 4-{1-Ethyl-1-[4-(2-hydroxy-2-isopropyl-3-methyl-butoxy)-3-methyl-phenyl]-propyl}-2-methyl-benzoic acid.



- Using a procedure analogous to Example 1I, 4-{1-ethyl-1-[4-(2-hydroxy-2-isopropyl-3-methyl-butoxy)-3-methyl-phenyl]-propyl}-2-methyl-benzoic acid methyl ester (0.30 g, 0.59 mmol) gives the title compound (0.96 g, 2.18 mmol, 56%).
- 15  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ),  $\delta$  0.63 (t,  $J = 7.3$  Hz, 6H), 1.00-1.06 (m, 12H), 2.10 (q,  $J = 7.2$  Hz, 4H), 2.12-2.19 (m, 2H), 2.17 (s, 3H), 2.62 (s, 3H), 3.89 (s, 2H), 6.71 (d,  $J = 8.4$  Hz, 1H), 6.86 (d,  $J = 1.7$  Hz, 1H), 6.94 (dd,  $J = 8.6, 2.5$  Hz, 1H), 7.08 (dd,  $J = 8.4, 1.7$  Hz, 1H), 7.09 (s, 1H), 7.94 (d,  $J = 8.4$  Hz, 1H).
- 20 ES-MS ( $m/z$ ): calcd. for  $\text{C}_{28}\text{H}_{39}\text{NO}_4$  (M-H): 439.6; found: 439.2.

I. Preparation of (4-{1-Ethyl-1-[4-(2-hydroxy-2-isopropyl-3-methyl-butoxy)-3-methyl-phenyl]-propyl}-2-methyl-benzoylamino)-acetic acid methyl ester.



-87-

Using a procedure analogous to Example 1J, 4-{1-ethyl-1-[4-(2-hydroxy-2-isopropyl-3-methyl-butoxy)-3-methyl-phenyl]-propyl}-2-methyl-benzoic acid (0.30 g, 0.68 mmol) and glycine methyl ester hydrochloride (0.094 g, 0.75 mmol) gives the titled compound (0.30 g, 0.59 mmol, 86%).

5 <sup>1</sup>H NMR (CDCl<sub>3</sub>), δ 0.62 (t, *J* = 7.5 Hz, 6H), 0.98-1.06 (m, 12H), 2.07 (q, *J* = 7.2 Hz, 4H), 2.10-2.19 (m, 2H), 2.16 (s, 3H), 2.44 (s, 3H), 3.80 (s, 2H), 3.89 (s, 1H), 4.23 (d, *J* = 5.1 Hz), 6.29 (t, *J* = 5.1 Hz), 6.70 (d, *J* = 8.7 Hz), 6.84 (d, *J* = 2.2 Hz, 1H), 6.94 (dd, *J* = 8.7, 2.2 Hz, 1H), 7.01 (d, *J* = 8.2 Hz, 1H), 7.05 (s, 1H), 7.32 (d, *J* = 8.2 Hz, 1H).

10 J. (4-{1-Ethyl-1-[4-(2-hydroxy-2-isopropyl-3-methyl-butoxy)-3-methyl-phenyl]-propyl}-2-methyl-benzoylamino)-acetic acid.

Using a procedure analogous to Example 1I, (4-{1-ethyl-1-[4-(2-hydroxy-2-isopropyl-3-methyl-butoxy)-3-methyl-phenyl]-propyl}-2-methyl-benzoylamino)-acetic acid methyl ester (0.30 g, 0.59 mmol) gives the title compound (0.25 g, 0.50 mmol, 86 %).

15 <sup>1</sup>H NMR (CDCl<sub>3</sub>), δ 0.61 (t, *J* = 7.4 Hz, 6H), 0.98-1.06 (m, 12H), 2.06 (q, *J* = 7.4 Hz, 4H), 2.10-2.19 (m, 2H), 2.16 (s, 3H), 2.43 (s, 3H), 3.89 (s, 2H), 4.25 (d, *J* = 5.2 Hz, 2H), 6.40 (t, *J* = 5.2 Hz, 1H), 6.70 (d, *J* = 8.6 Hz, 1H), 6.83 (d, *J* = 2.0 Hz, 1H), 6.93 (dd, *J* = 8.3, 2.0 Hz, 1H), 7.01 (d, *J* = 8.0 Hz, 1H), 7.05 (s, 1H), 7.31 (d, *J* = 8.0 Hz, 1H).

20 ES-MS (*m/z*): calcd. for C<sub>30</sub>H<sub>42</sub>NO<sub>5</sub> (M-H)<sup>-</sup>: 496.3; found: 496.6.

#### Compounds of the Invention – Salts, Stereoisomers, & Prodrugs:

25 Salts of the compounds represented by formulae I are an additional aspect of the invention. The skilled artisan will also appreciate that the family of compounds of formulae I include acidic and basic members and that the present invention includes pharmaceutically acceptable salts thereof.

In those instances where the compounds of the invention possess acidic or basic functional groups various salts may be formed which are more water soluble and physiologically suitable than the parent compound. Representative pharmaceutically acceptable salts, include but are not limited to, the alkali and alkaline earth salts such as

30



-88-

lithium, sodium, potassium, ammonium, calcium, magnesium, aluminum, zinc, and the like. Sodium and potassium salts are particularly preferred. Salts are conveniently prepared from the free acid by treating the acid in solution with a base or by exposing the acid to an ion exchange resin. For example, a carboxylic acid substituent on the  
5 compound of Formula I may be selected as  $-CO_2H$  and salts may be formed by reaction with appropriate bases (e.g., NaOH, KOH) to yield the corresponding sodium and potassium salt.

Included within the definition of pharmaceutically acceptable salts are the relatively non-toxic, inorganic and organic base addition salts of compounds of the  
10 present invention, for example, ammonium, quaternary ammonium, and amine cations, derived from nitrogenous bases of sufficient basicity to form salts with the compounds of this invention (see, for example, S. M. Berge, *et al.*, "Pharmaceutical Salts," J. Phar. Sci., 66: 1-19 (1977)). Moreover, the basic group(s) of the compound of the invention may be reacted with suitable organic or inorganic acids to form salts such as acetate,  
15 benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, camsylate, carbonate, chloride, choline, clavulanate, citrate, chloride, chloroprocaine, choline, diethanolamine, dihydrochloride, diphosphate, edetate, edisylate, estolate, esylate, ethylenediamine, fluoride, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, bromide, chloride, hydrobromide, hydrochloride,  
20 hydroxynaphthoate, iodide, isothionate, lactate, lactobionate, laurate, malate, maleate, malseate, mandelate, meglumine, mesylate, mesviate, methylbromide, methylnitrate, methylsulfate, mucate, napsylate, nitrate, oleate, oxalate, palmitate, pamoate, pantothenate, phosphate, polygalacturonate, procaine, salicylate, stearate, subacetate, succinate, sulfate, tannate, tartrate, teoclate, tosylate, trifluoroacetate, trifluoromethane  
25 sulfonate, and valerate.

Certain compounds of the invention may possess one or more chiral centers and may thus exist in optically active forms. Likewise, when the compounds contain an alkenyl or alkenylene group there exists the possibility of cis- and trans- isomeric forms of the compounds. The R- and S- isomers and mixtures thereof, including racemic mixtures  
30 as well as mixtures of cis- and trans- isomers, are contemplated by this invention. Additional asymmetric carbon atoms can be present in a substituent group such as an alkyl group. All such isomers as well as the mixtures thereof are intended to be included

-89-

in the invention. If a particular stereoisomer is desired, it can be prepared by methods well known in the art by using stereospecific reactions with starting materials which contain the asymmetric centers and are already resolved or, alternatively by methods which lead to mixtures of the stereoisomers and subsequent resolution by known methods. For example, a chiral column may be used such as those sold by Daicel Chemical Industries identified by the trademarks:

CHIRALPAK AD, CHIRALPAK AS, CHIRALPAK OD, CHIRALPAK OJ,  
CHIRALPAK OA, CHIRALPAK OB, CHIRALPAK OC, CHIRALPAK OF,  
CHIRALPAK OG, CHIRALPAK OK, and  
CHIRALPAK CA-1.

By another conventional method, a racemic mixture may be reacted with a single enantiomer of some other compound. This changes the racemic form into a mixture of diastereomers. These diastereomers, because they have different melting points, different boiling points, and different solubilities can be separated by conventional means, such as crystallization.

The present invention is also embodied in mixtures of compounds of formulae I.


Prodrugs are derivatives of the compounds of the invention which have chemically or metabolically cleavable groups and become by solvolysis or under physiological conditions the compounds of the invention which are pharmaceutically active in vivo.

Derivatives of the compounds of this invention have activity in both their acid and base derivative forms, but the acid derivative form often offers advantages of solubility, tissue compatibility, or delayed release in a mammalian organism (see, Bundgard, H., Design of Prodrugs, pp. 7-9, 21-24, Elsevier, Amsterdam 1985). Prodrugs include acid derivatives well known to practitioners of the art, such as, for example, esters prepared by reaction of the parent acidic compound with a suitable alcohol, or amides prepared by reaction of the parent acid compound with a suitable amine. Simple aliphatic or aromatic esters derived from acidic groups pendent on the compounds of this invention are preferred prodrugs. In some cases it is desirable to prepare double ester type prodrugs such as (acyloxy) alkyl esters or ((alkoxycarbonyl)oxy)alkyl esters. Particularly preferred esters to use as prodrugs are; methyl, ethyl, propyl, isopropyl, n-butyl, isobutyl, morpholinoethyl, and N,N-diethylglycolamido.


-90-

N,N-diethylglycolamido ester prodrugs may be prepared by reaction of the sodium salt of a compound of Formula (I) (in a medium such as dimethylformamide) with 2-chloro-N,N-diethylacetamide (available from Aldrich Chemical Co., Milwaukee, Wisconsin USA; Item No.25,099-6).

- 5 Morpholinylethyl ester prodrugs may be prepared by reaction of the sodium salt of a compound of Formula I (in a medium such as dimethylformamide) 4-(2-chloroethyl)morpholine hydrochloride (available from Aldrich Chemical Co., Milwaukee, Wisconsin USA, Item No. C5,220-3). For example, prodrugs may be prepared by reaction of the sodium salt for a compound of Formula I with;




and sodium iodide to provide the ester prodrug pendent group




Also, lower alkyl (viz., C<sub>1</sub>-C<sub>8</sub>) ester prodrugs may be prepared by conventional means such as reacting the sodium or potassium salt (derived by forming the salt of any acidic compound of the invention; viz., reaction of a base such as KOH with an acidic group such as -CO<sub>2</sub>H) of a compound of Formula I with an alkyl iodide such as methyl iodide, ethyl iodide, n-propyl iodide, isopropyl iodide. Typical ester prodrug substituents are

-91-



or



Pharmaceutical Formulations containing the Novel Compounds of the Invention:

Pharmaceutical formulations of the invention are prepared by combining (e.g.,  
5 mixing) a therapeutically effective amount of the compound of the invention  
(compounds of Formula I ) together with a pharmaceutically acceptable carrier or  
diluent. The present pharmaceutical formulations are prepared by known procedures  
using well-known and readily available ingredients.

In making the compositions of the present invention, the compounds of  
10 Formula I will usually be admixed with a carrier, or diluted by a carrier, or enclosed  
within a carrier which may be in the form of a capsule, sachet, paper or other container.  
When the carrier serves as a diluent, it may be a solid, semi-solid or liquid material  
which acts as a vehicle, or can be in the form of tablets, pills, powders, lozenges,  
elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid  
15 medium), or ointment, containing, for example, up to 10% by weight of the  
compound. The compounds of the present invention are preferably formulated prior to  
administration.

The compounds of the invention may also be delivered by suitable formulations  
contained in a transderm patch. Alternatively, the compounds of the invention may be  
20 delived to a patient by sublingual administration.

For the pharmaceutical formulations any suitable carrier known in the art can be  
used. In such a formulation, the carrier may be a solid, liquid, or mixture of a solid and a  
liquid. Solid form formulations include powders, tablets and capsules. A solid carrier  
can be one or more substances which may also act as flavoring agents, lubricants,  
25 solubilisers, suspending agents, binders, tablet disintegrating agents and encapsulating  
material.

Tablets for oral administration may contain suitable excipients such as calcium  
carbonate, sodium carbonate, lactose, calcium phosphate, together with disintegrating  
agents, such as maize, starch, or alginic acid, and/or binding agents, for example,

-92-

gelatin or acacia, and lubricating agents such as magnesium stearate, stearic acid, or talc.

In powders the carrier is a finely divided solid which is in admixture with the finely divided Active ingredient. In tablets the compound of Formula I is mixed with a carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain from about 1 to about 99 weight percent of the compound which is the novel compound of this invention. Suitable solid carriers are magnesium carbonate, magnesium stearate, talc, sugar lactose, pectin, dextrin, starch, gelatin, tragacanth, methyl cellulose, sodium carboxymethyl cellulose, low melting waxes, and cocoa butter.

Sterile liquid form formulations include suspensions, emulsions, syrups and elixirs.

The Active Ingredient may be dissolved or suspended in a pharmaceutically acceptable carrier, such as sterile water, sterile organic solvent or a mixture of both. The compounds can often be dissolved in a suitable organic solvent, for instance aqueous propylene glycol. Other compositions can be made by dispersing the finely divided compounds of the invention in aqueous starch or sodium carboxymethyl cellulose solution or in a suitable oil.

#### Methods of Using the Compounds of the Invention:

Many disease states are benefited by treatment with the compounds of Formula I include, but are not limited to:

- disease states characterized by abnormal calcium regulation
- disease states characterized by abnormal cell proliferation
- disease states characterized by abnormal cell differentiation
- disease states characterized by abnormal immune response
- disease states characterized by abnormal dermatological conditions
- disease states characterized by neurodegenerative condition
- disease states characterized by inflammation
- disease states characterized by vitamin D sensitivity
- disease states characterized by hyperproliferative disorders.

-93-

Specific disease states benefited by treatment of the compounds of Formula I and II include, but are not limited to:

|    |                                  |
|----|----------------------------------|
|    | Acne                             |
|    | Actinic keratosis                |
| 5  | Alopecia                         |
|    | Alzheimer's disease              |
|    | Benign prostatic hyperplasia     |
|    | Bladder cancer                   |
|    | Bone maintenance in zero gravity |
| 10 | Bone fracture healing            |
|    | Breast cancer                    |
|    | Chemoprevention of Cancer        |
|    | Crohn's disease                  |
|    | Colon cancer                     |
| 15 | Type I diabetes                  |
|    | Host-graft rejection             |
|    | Hypercalcemia                    |
|    | Type II diabetes                 |
|    | Leukemia                         |
| 20 | Multiple sclerosis               |
|    | Myelodysplastic syndrome         |
|    | Insufficient sebum secretion     |
|    | Osteomalacia                     |
|    | Osteoporosis                     |
| 25 | Insufficient dermal firmness     |
|    | Insufficient dermal hydration    |
|    | Psoriatic arthritis              |
|    | Prostate cancer                  |
|    | Psoriasis                        |
| 30 | Renal osteodystrophy             |
|    | Rheumatoid arthritis             |
|    | Scleroderma                      |

-94-

Skin cancer

Systemic lupus erythematosus

Skin cell protection from Mustard vesicants

Ulcerative colitis

5

Vitiligo

Wrinkles

Particularly preferred is the treatment of psoriasis and osteoporosis by administration to a mammal (including a human) of a therapeutically effective amount of compounds of Formulae I. By "pharmaceutically effective amount" it is meant that  
10 quantity of pharmaceutical agent corresponding to formulae I which prevents, removes or reduces the deleterious effects of a disease state in mammals, including humans.

The specific dose of a compound administered according to this invention to obtain therapeutic or prophylactic effects will, of course, be determined by the particular circumstances surrounding the case, including, for example, the compound administered,  
15 the route of administration and the condition being treated. Typical daily doses will contain a pharmaceutically effective amount typically in the range of from about 0.0001 mg/kg/day to about 50 mg/kg/day of body weight of an active compound of this invention. Preferably the dose of compounds of the invention will be from 0.0001 to 5 mg/kg/day of body weight.

20 Preferably compounds of the invention (e.g., per Formula I) or pharmaceutical formulations containing these compounds are in unit dosage form for administration to a mammal. The unit dosage form can be a capsule or tablet itself, or the appropriate number of any of these. The quantity of Active ingredient in a unit dose of composition may be varied or adjusted from about 0.0001 to about 1000 milligrams or more  
25 according to the particular treatment involved. It may be appreciated that it is necessary to make routine variations to the dosage depending on the age and condition of the patient. Dosage will also depend on the route of administration. The compounds of the invention may be administered by a variety of routes including oral, aerosol, rectal, transdermal, sublingual, subcutaneous, intravenous, intramuscular, and  
30 intranasal. Particularly preferred is the treatment of psoriasis with an ointment type formulation containing the compounds of the invention. The ointment formulation may be applied as needed, typically from one to 6 times daily.

-95-

Treatment of psoriasis is preferably done with topical application by a formulation in the form of a cream, oil, emulsion, paste or ointment containing a therapeutically effective amount of a compound of the invention.. The formulation for topical treatment contains from 0.5 to 0.00005 weight percent, preferably from .05 to 5 0.0005 weight percent, and most preferably from 0.025 to 0.001 of a Active Ingredient.

For example, two semisolid topical preparations useful as vehicles for VDR modulators in treatment and prevention of psoriasis are as follows:

Polyethylene Glycol Ointment USP (p. 2495)

Prepare Polyethylene Glycol Ointment as follows:

|    |                          |               |
|----|--------------------------|---------------|
| 10 | Polyethylene Glycol 3350 | 400 g.        |
|    | Polyethylene Glycol 400  | <u>600 g.</u> |
|    | To make                  | 1000 g.       |

Heat the two ingredients on a water bath to 65C. Allow to cool, and stir until congealed. If a firmer preparation is desired, replace up to 100 g of 15 the polyethylene glycol 400 with an equal amount of polyethylene glycol 3350.

Hydrophilic Ointment USP (p. 1216)

Prepare Hydrophilic Ointment as follows:

|    |                       |               |
|----|-----------------------|---------------|
|    | Methylparaben         | 0.25 g.       |
| 20 | Propylparaben         | 0.15 g.       |
|    | Sodium Lauryl Sulfate | 10 g.         |
|    | Propylene Glycol      | 120 g.        |
|    | Stearyl Alcohol       | 250 g.        |
|    | White Petrolatum      | 250 g.        |
| 25 | Purified Water        | <u>370 g.</u> |
|    | To make about         | 1000 g.       |

The Stearyl Alcohol and White Petrolatum are melted on a steam bath, and warmed to about 75C. The other ingredients, previously dissolved in the water are added, warmed to 75C, and the mixture stirred until it congeals.

30 For each of the above formulations the Active Ingredient is added during the heating step in an amount that is from 0.5 to 0.00005 weight percent, preferably from .05 to 0.0005 weight percent, and most preferably from 0.025 to 0.001 weight percent of the



-96-

total ointment weight. (Source: - United States Pharmacopoeia 24, United States Pharmacopeial Convention, 1999)

Conventional therapy for osteoporosis includes; (i) estrogens, (ii) androgens, (iii) calcium supplements, (iv) vitamin D metabolites, (v) thiazide diuretics, (vi) calcitonin, (vii) bisphosphonates, (viii) SERMS, and (ix) fluorides (see, Harrison's Principles of Internal Medicine, 13<sup>th</sup> edition, 1994, published by McGraw Hill Publ., ISBN 0-07-032370-4, pgs.2172-77; the disclosure of which is incorporated herein by reference.). Any one or combination of these conventional therapies may be used in combination with the method of treatment using compounds of Formulae I as taught herein. For example, in a method of treating osteoporosis, the vitamin D receptor modulator compounds of the invention (e.g., as defined by formula I) may be administered separately or simultaneously with a conventional therapy. Alternatively, the vitamin D receptor modulator compounds of the invention may be combined with conventional therapeutic agents in a formulation for treatment of osteoporosis such as set out below:

A formulation for treating osteoporosis comprising:

Ingredient (A1): a vitamin D receptor modulator represented by formula (I), or a pharmaceutically acceptable salt or prodrug derivative thereof;

Ingredient (B1):

one or more co-agents that are conventional for treatment osteoporosis selected from the group consisting of:

- a. estrogens,
- b. androgens,
- c. calcium supplements,
- d. vitamin D metabolites,
- e. thiazide diuretics,
- f. calcitonin,
- g. bisphosphonates,
- h. SERMS, and
- i. fluorides.

Ingredient (C1): optionally, a carrier or diluent.

-97-

Typically useful formulations are those wherein the weight ratio of (A1) to (B1) is from 10:1 to 1:1000 and preferably from 1:1 to 1:100.

#### Combination Therapy for Psoriasis:

5           Conventional therapy for psoriasis includes topical glucocorticoids, salicylic acid, crude coal tar, ultraviolet light, and methotrexate (see, Harrison's Principles of Internal Medicine, 13<sup>th</sup> edition, 1994, published by McGraw Hill Publ., ISBN 0-07-032370-4, pgs.2172-77). Any one or combination of these conventional therapies may be used in combination with the method of treatment using compounds of Formulae I as taught  
10       herein. For example, in a method of treating osteoporosis, the vitamin D receptor modulator compounds of the invention (e.g., as defined by formula I) may be topically administered separately or simultaneously with a conventional therapy. Alternatively, the vitamin D receptor modulator compounds of the invention may be combined with conventional therapeutic agents in a topically applied formulation for treatment of  
15       osteoporosis such as set out below:

A formulation for treating osteoporosis comprising:

Ingredient (A2):       a vitamin D receptor modulator represented by formula (I), or a pharmaceutically acceptable salt or prodrug derivative thereof;

20           Ingredient (B2):  
                  one or more co-agents that are conventional for treatment of osteoporosis selected from the group consisting of:

- a.       topical glucocorticoids,  
                  b.       salicylic acid, or  
25               c.       crude coal tar.

Ingredient (C2): optionally, a carrier or diluent.

Typically useful formulations are those wherein the weight ratio of (A2) to (B2) is from 1:10 to 1:100000 and preferably from 1:100 to 1:10000.

30

#### Experimental Results:

Table 1 of Experimental Results

-98-

| Test<br>Cmpd. <sup>1</sup> | RXR-VDR<br>(SaOS-2<br>cells) <sup>2</sup><br>EC <sub>50</sub> (nM) | VDR CTF<br>(Caco-2 cells) <sup>3</sup><br>EC <sub>50</sub> (nM) | OCN<br>Promoter <sup>4</sup><br>EC <sub>50</sub> (nM) | Mouse<br>Hypercal <sup>5</sup><br>μg/Kg/d |
|----------------------------|--|---|---|---|
| Ex. 1                      | 68.6   | 471   | 0.96  | 3000                                      |
| Ex. 2                      | 111.9/9.4  | 397   | 1.2/1.7   | 300                                       |
| Ex. 3                      | 300.7/148.9  | 819   | 16.3/19.7   | 1000                                      |
| Ex. 4                      | 257/127.3  | 974   | 21/27.7   | 73000                                     |
| AA                         | 5.02   | 16  | 5   | 0.06                                      |
| BB                         | 10.32  | 169.81  | 8.24  | 20  |
| CC                         | 2427.7   |   | 2680.9  |   |
| DD                         | 109.44   |   | 31.1  | 1000                                      |
| EE                         | 429.99   | 891.16  | 341.25  | 1000                                      |
| FF                         | 3/1  | 57  | 0.28  |   |

Table 2 of Experimental Results

| Test<br>Cmpd. <sup>1</sup> | Kera. Prolif.<br>IC <sub>50</sub> (nM) | IL-10<br>IC <sub>50</sub> (nM) |
|----------------------------|--|--------------------------------|
| Ex. 1                      |  |                                |
| Ex. 2                      | 21                                     | 24                             |
| Ex. 3                      | 121                                    | 718                            |

-99-

|       |      |     |
|-------|------|-----|
| Ex. 4 |      | 477 |
| AA    | 120  | 1.2 |
| BB    | 10   | 28  |
| CC    |      |     |
| DD    | 1060 |     |
| EE    |      |     |
| FF    | 103  | 0.5 |


Explanation of Table 5 and 6 column numerical superscripts:

1. Test Compound numbers refer to the products of the corresponding Example Nos. that is, compounds within the scope of the invention. For example, the number "Ex. 2" refers to the compound, 3'-[4-(2-hydroxy-3,3-dimethylbutoxy)-3-methylphenyl]-3'-[5-methoxycarbonyl-4-methylthiophen-2-yl]pentane, prepared in Example 2. The control experiments are done with the double letter coded compounds identified as follows:

"AA" = 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub>

"BB" = 3-(4-{1-Ethyl-1-[4-(2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-2-methyl-phenoxy)-propane-1,2-diol


"CC" = 1-(4-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-cyclohexyl}-2-methyl-phenoxy)-3,3-dimethyl-butan-2-one




"DD" = compound represented by the formula:

"EE" = compound represented by the formula:

-100-



“FF” = calcipotriol (structural formula below):



2. The RXR-VDR heterodimerization (SaOS-2 cells) test is described in the “Assay” section of the Description, *infra*.

3. The VDR CTF (Caco-2 cells) test is described in the “Assay” section of the Description, *infra*.

4. The OCN Promoter test is described in the “Assay” section of the Description, *infra*.

5. The Mouse Hypercalcemia test is described in the “Assay” section of the Description, *infra*.

6. The keratinocyte proliferation assay is described in the “Assay” section of the Description, *infra*.

7. The IL-10 induction assay is described in the “Assay” section of the Description, *infra*.

## Assay Methods

### Use of the Assay Methods:

The evaluation of the novel compounds of the invention for osteoporosis and other related diseases is done using a plurality of test results. The use of multiple assays is necessary since the combined properties of (i) high activity for the vitamin D receptor, and (ii) prevention of hypercalcemia must be achieved to have utility for the methods of treating diseases, which are also, aspects of this invention. Some of the tests described

-101-

below are believed related to other tests and measure related properties of compounds. Consequently, a compound may be considered to have utility in the practice of the invention if it meets most, if not all, of the acceptance criteria for the above described tests.

- 5           The evaluation of the novel compounds of the invention for psoriasis is done using the Keratinocyte Proliferation Assay in combination with other assays that measure inhibition of IL-2 production and stimulation of IL-10 production in peripheral blood mononuclear cells (PBMCs).

Brief Description, Utility and Acceptance Criteria for the Assay Methods:

10

1. The RXR-VDR heterodimer Assay:

- This assay provides the VDR activity of a test compound. It is desirable to have low EC50 values for a compound in this assay. The lower the EC50 value, the more active the compound will be as a VDR agonist. Desired assay results are EC50 values less than or equal to 600 nM. Preferred assay results are less than 250 nM, and most preferably less than 150 nM.

15

2. The Caco-2 cell Co-transfection Assay:

- The Caco-2 cell assay is an indicator for the undesirable condition of hypercalcemia. This co-transfection assay is a surrogate assay for in vivo calcemic activity of VDR ligands. It is desirable to have high EC50 values for a test compound in this assay. The higher the EC50 values for a compound the less calcemic it will be in vivo. Desired assay results are EC50 greater than or equal to 300 nM. Preferred assay results are greater than 1000 nM.

20

3. The OCN (osteocalcin) Promoter Assay

- The OCN Promoter Assay is an indicator and marker for osteoporosis. Desired assay results are EC50 less than or equal to 325 nM. Preferred assay results are less than 50 nM.

25

4. The Mouse Hypercalcemia Assay

30

-102-

The Mouse Hypercalcemia Assay is a six day hypercalcemia test for toxicity and selectivity. Acceptable test results are levels greater than 300 µg/kg/day. Preferred assay results are levels greater than 1000 µg/kg/day.

5    5. The Keratinocyte Proliferation Assay

This Assay is indicative for the treatment of psoriasis. An acceptable test result is IC<sub>50</sub> value of less than or equal to 300 nM. Preferred assay results are IC<sub>50</sub> values of less than 100 nM.

10   6. The IL-10 induction Assay

This is an in vitro efficacy assay for psoriasis, abscess and adhesion. Psoriasis involves both keratinocytes and immune cells. IL-10 is a unique cytokine because it is anti-inflammatory and immunosuppressive. This assay tells us whether a VDRM is able to function as an agonist in PBMCs (primary blood mononuclear cells) or not. A lower EC<sub>50</sub> value is desirable in this assay since a compound with a lower EC<sub>50</sub> value will be a better agonist in PBMCs. An acceptable test result is an EC<sub>50</sub> value of less than 200 nM. Preferred assay results are EC<sub>50</sub> values of less than 100 nM.

7. Other Compound Assay Standards

20   An alternative measure of the efficacy of compounds of the invention for treatment of osteoporosis is a numerical ratio calculated as follows:

$$\frac{\text{Dose Threshold needed to induce hypercalcemia}}{\text{divided by}} \\ \text{Dose Threshold needed for bone efficacy}$$

25

An alternative measure of the efficacy of compounds of the invention for treatment of psoriasis is a numerical ratio calculated as follows:

$$\frac{\text{Dose Threshold needed to induce hypercalcemia}}{\text{divided by}}$$

30

$$\text{Dose Threshold needed to induce keratinocyte proliferation}$$

For the above ratios, Dose Thresholds are determined from dose response curve data.

-103-

8. The CaT1 (calcium transport protein 1) Assay

The CaT1 Assay is an indicator for the undesirable condition of hypercalcemia. The higher the EC50 values for a compound the less calcemic it will be in vivo. Desired assay results are EC50 greater than or equal to 500 nM. Preferred assay results are greater than 1000 nM.

Details of the Assay Methods: . . . .(1) Materials and Method for RXR-VDR Heterodimerization Assay:

## Transfection Method:

- 10 • FuGENE 6 Transfection Reagent (Roche Cat # 1 814 443 )

## Growth Media:

- D-MEM High Glucose (Gibco BRL Cat # 11054-020), 10% FBS, 1% antibiotic-antimycotic (Ab-Am)

FBS heat inactivated (Gibco BRL Cat # 10092-147 )

- 15 Ab-Am (Gibco BRL Cat # 15240-062 )

## Cells:

- Grow SaOs-2 cells in T-152 cm<sup>2</sup> culture flasks in *growth media*.
- Keep the density at 5-6 x 10<sup>5</sup> cells/ml
- Passage cells 1:3 twice a week
- 20 • Add Trypsin EDTA (Gibco BRL Cat # 25300-020) and incubate
- Resuspend cells in plating media and transfer into growth media.

## Wash Media:

- HBSS Low Glucose Without Phenol Red (Gibco BRL Cat # 14175-095), 1% Ab-Am

## Plating Media:

- 25 • D-MEM Low Glucose Without Phenol Red (Gibco BRL Cat # 11054-020), 1% Ab-Am
- D-MEM

Stripped FBS (Hyclone Cat# SH30068.03 Lot # AHM9371 )

Ab-Am

## Transfection / Treatment Media:

- 30 • D-MEM Low Glucose Without Phenol Red only

T-152 cm<sup>2</sup> culture flask:

- Use Corning Costar T-152 cm<sup>2</sup> culture flask (Cat # 430825) to grow the cells



-104-

Flat well Plates:

- Use well plate to plate cells
- Use Deep well plate sterile to make up treatment media.

5 Luciferase Assay Reagent:

- Use Steady-Glo Luciferase Reagent from Promega (Cat # E2550) Consists of:
  - a. E2533 Assay Substrate, lyophilized product and
  - b. E2543 Assay Buffer.
- Thaw at room temperature

10 • Store

DAY 1: Cell Plating:

Cell Harvesting

Aspirate media from culture flask, rinse cells with HBSS and aspirate.

Add trypsin and incubate.

15 When cells appear detached, resuspend cells in *growth media*.

Transfer into a new flask with fresh *growth media* for passaging the cells.

Plate well plates and two extra plates

A. Cell Count

Mix the cell suspension using pipette

20 Use *Hemocytometer* to count the cells

Load cell suspension onto the hemocytometer chamber

Count cells.

Plate seeding:

Use plating media 10 % Stripped FBS in D-MEM Low Glucose, Without Phenol Red, 1%

25 Ab-Am

Plate 14 plates @ 165  $\mu$ l / well.

In sterile flask add cell suspension

to *plating media*.

Mix .

30 Add cells / well.

Place the cells in the incubator.

Cells should be about 75 % confluent prior to transfection.

-105-

## Step 1: DNA and Media

Add plain DMEM media to tubes for mixing the DNA

Add the Reporter gene pFR-LUC

- 5 Add the Gal4-RXR-DEF and VP16-VDR-LBD

## Step 2: FuGENE and Media

Prepare plain DMEM media in a tubes for mixing FuGENE

Add *FuGENE 6 Transfection Reagent*

- 10 Incubate

## Step 3: FuGENE , DNA and Media Complex

Add FuGENE Media complex from step 2 to DNA Media complex from step1

Incubate

15

## Step 4: FuGENE , DNA and Media Complex to-well plate

Add FuGENE-DNA-Media complex from step 3 to each plate

Incubate.

20

## Day 3: Dosing

Treatment preparation

Allow for transfection time

Make a stock solution of the compounds in DMSO

Vortex until all the compounds has been dissolved.

- 25 Further dilute in D-MEM (Low Glucose – With out Phenol Red)

Add compounds in quadruplicate to give final volume

Incubate.

Day 4: Luciferase Assay

Read the plates after drug treatment

- 30 Remove part of media from all the wells and leave remainder

Add Steady-Glo Luciferase Reagent mixture / wells

Incubate

-106-

Count each well using a Luminescence counter, Top Count NXT by Packard

Set a delay between plates to reduce the background.

(2) Materials and Method for The Caco-2 Cell Assay:

5 Caco-2 cells, grown in phenol red free, DMEM (Invitrogen, Carlsbad, CA) containing 10 % charcoal-stripped FCS (Hyclone, Logan, UT), were transfected with Fugene 6 reagent (Roche Diagnostics, Indianapolis, IN). Cells (5000/well) were plated 18 h before transfection in a 96 well plate. The Cells were transfected with Gal4-responsive reporter pFRLuc (150 ng, Stratagene, La Jolla CA) and the  
10 receptor expression vector pGal4-VDR-LBD (10 ng), along with Fugene 6 reagent (0.2  $\mu$ l/well). The DNA-Fugene complex was formed by incubating the mixture for 30 min at room temperature. The cells were transfected in triplicate for 5 h, and treated with various concentrations of VDR ligands (from 0.01 nM to 10,000 nM concentration range) 18h post-transfection. The luciferase activity  
15 was quantified using Steady-Glo reagent kit (Promega, Madison, WI) as per manufacturer's specifications.

(3) Materials and Method for The OCN Promoter Assay:

20 The activation of osteocalcin by VDR ligands was evaluated in a rat osteoblast-like cell line RG-15 (ROS 17/2.8) stably expressing rat osteocalcin promoter fused with luciferase reporter gene. The stable cell lines were established as reported before (Activation of Osteocalcin Transcription involves interaction of protein kinase A- and Protein kinase C-dependent pathways.  
25 Boguslawski, G., Hale, L. V., Yu, X.-P., Miles, R. R., Onyia, J. E., Santerre R. F., Chandrasekhar, S. J Biol. Chem. 275, 999-1006, 2000). Confluent RG-15 cells maintained in DMEM/F-12 medium (3:1) containing 5% FBS, 300  $\mu$ g/ml G418 and at 37°C under 5% CO<sub>2</sub>/95% air atmosphere were trypsinized (0.25% trypsin) and plated into white opaque 96-well cell culture plates (25000  
30 cells/well). After 24 hr, cells (in DMEM/F-12 medium + 2% FBS) were treated with various concentrations of compounds, dissolved in DMSO. The final DMSO concentration remained at 0.01% (v/v). After 48 hr treatment, the

-107-

medium was removed, cells were lysed with 50  $\mu$ l of lysis buffer (From Luciferase reporter assay system, Roche Diagnostics, Indianapolis, IN) and assayed for luciferase activity using the Luciferase Reporter Gene Assay kit from Boehringer Mannheim as per manufacturer's specifications.

5

(4) Materials and Method for The Mouse Hypercalcemia Assay:

Weanling, virus -antibody-free, five to six weeks old female DBF mice (Harlan, Indianapolis, IN) are used for all the studies. Animals are allowed to acclimate to local vivarium conditions for 2 days. Mice are maintained on a 12 hr light/dark cycle at 22<sup>0</sup>C with ad lib access to food (TD 5001 with 1.2% Ca and 0.9%P, Teklad, Madison, WI) and water. The animals then are divided into groups with 4-5 mice per group. Different doses of test compounds prepared in 10% Ethanol and 90% sesame oil are administered to mice orally via gavage for 6 days. 1 $\alpha$ -25(OH)<sub>2</sub>D<sub>3</sub> 0.5 $\mu$ g/kg/d was also given to one group of mice as the positive control. Serum ionized calcium is evaluated at 6 hours after the last dosing under isoflurane anesthesia by Ciba-Corning Ca<sup>++</sup>/PH Analyzer, (Model 634, Chiron Diagnostics Corp., East Walpole, MA). Raw data of group differences is assessed by analysis of variance (ANOVA) using Fisher's protected least significant difference (PLSD) where the significance level was P< 0.05.

20 (5) The Keratinocyte Proliferation Assay:

KERtr cells (Human skin keratinocyte transformed with a retrovirus vector, obtained from ATCC) were plated in 96-well flat-bottomed plates (3000 cells/well) in 100  $\mu$ l keratinocyte serum free medium supplemented with bovine pituitary extract in the absence of EGF (Life Technologies, Rockville, MD) and incubated at 37°C for two days. The cells were treated with various concentrations of VDR ligands (ten-fold serial dilution from 10,000 nM to 0.1 nM in triplicate), dissolved in 100  $\mu$ l keratinocyte serum free medium supplemented with bovine pituitary extract in the absence of EGF and incubated at 37°C for 72hr. BrdU (5-bromo-2'-deoxyuridine) incorporation was analyzed as a measure of DNA replication (Cell proliferation ELISA kit, Roche Diagnostics, Indianapolis, IN) and absorbance was measured at 405 nm. Potency values (IC<sub>50</sub>) values were determined as the concentration (nM) of compound that elicited a half-maximal response.

30

(6) Materials and Method for human IL-10 Induction Assay:

-109-

serum (Invitrogen, Carlsbad, CA), are plated at 5500 cell per well in a 96-well plate in a total volume of 100  $\mu$ l/well. The cells are kept in the 96-well plate for 6 days to differentiate them to small intestinal cells that express the calcium transporter, CaT1. On day 3 after plating, old media is removed and replaced with fresh media (150  $\mu$ l/well). On day 6 the old media is removed and the cells are kept in treatment media (180  $\mu$ l/well) that contained 10 % charcoal stripped fetal bovine serum (Hyclone, Logan, UT) in DMEM (low glucose, without phenol red; Invitrogen, Carlsbad, CA). The cells are treated with various concentrations of VDR ligands (from 0.01 nM to 10,000 nM concentration range) prepared in treatment media (20  $\mu$ l/well). Twenty hours post-treatment, total RNA is prepared by RNeasy 96 method as described by the manufacturer (Qiagen, Valencia, CA). The RNA is reverse transcribed and amplified for human CaT1 and GAPDH (control) messages by quantitative RT-PCR using ABI PRISM 7900HT Sequence Detection System according to manufacturer's instructions (Applied Biosystems, Foster City, CA). Optimized primer pairs and probes for human CaT1 and GAPDH genes are obtained commercially (Applied Biosystems, Foster City, CA). Each 20  $\mu$ l quantitative RT-PCR reaction in a 384-well Taqman PCR plate consists of forward and reverse primers (900 nM), Taqman probe (200 nM), total RNA (4  $\mu$ l from each well of the 96-well culture plate) and 10  $\mu$ l of Taqman Universal PCR Master Mix (Roche Diagnostics, Indianapolis, IN). Reactions are incubated at 48°C for 30 minutes, followed by 10 minutes at 95°C and subjected to 40 cycles of PCR (95°C for 15 seconds followed by 60°C for 1 minute). GAPDH is used as an internal control and its primer and probe set are obtained commercially (Applied Biosystems, Foster City, CA).

-102-

The Mouse Hypercalcemia Assay is a six day hypercalcemia test for toxicity and selectivity. Acceptable test results are levels greater than 300 µg/kg/day. Preferred assay results are levels greater than 1000 µg/kg/day.

5     5. The Keratinocyte Proliferation Assay

      This Assay is indicative for the treatment of psoriasis. An acceptable test result is IC50 value of less than or equal to 300 nM. Preferred assay results are IC50 values of less than 100 nM.

10    6. The IL-10 induction Assay

      This is an in vitro efficacy assay for psoriasis, abscess and adhesion. Psoriasis involves both keratinocytes and immune cells. IL-10 is a unique cytokine because it is anti-inflammatory and immunosuppressive. This assay tells us whether a VDRM is able to function as an agonist in PBMCs (primary blood mononuclear cells) or not. A lower  
15    EC50 value is desirable in this assay since a compound with a lower EC50 value will be a better agonist in PBMCs. An acceptable test result is an EC50 value of less than 200 nM. Preferred assay results are EC50 values of less than 100 nM.

7. Other Compound Assay Standards

20    An alternative measure of the efficacy of compounds of the invention for treatment of osteoporosis is a numerical ratio calculated as follows:

$$\frac{\text{Dose Threshold needed to induce hypercalcemia}}{\text{Dose Threshold needed for bone efficacy}}$$

25

An alternative measure of the efficacy of compounds of the invention for treatment of psoriasis is a numerical ratio calculated as follows:

$$\frac{\text{Dose Threshold needed to induce hypercalcemia}}{\text{Dose Threshold needed to induce keratinocyte proliferation}}$$

30

For the above ratios, Dose Thresholds are determined from dose response curve data.

-103-

8. The CaT1 (calcium transport protein 1) Assay

The CaT1 Assay is an indicator for the undesirable condition of hypercalcemia. The higher the EC50 values for a compound the less calcemic it will be in vivo. Desired assay results are EC50 greater than or equal to 500 nM. Preferred assay results are greater than 1000 nM.

Details of the Assay Methods:(1) Materials and Method for RXR-VDR Heterodimerization Assay:

## Transfection Method:

- 10 • FuGENE 6 Transfection Reagent (Roche Cat # 1 814 443 )

## Growth Media:

- D-MEM High Glucose (Gibco BRL Cat # 11054-020), 10% FBS, 1% antibiotic-antimycotic (Ab-Am)

FBS heat inactivated (Gibco BRL Cat # 10092-147 )

- 15 Ab-Am (Gibco BRL Cat # 15240-062 )

## Cells:

- Grow SaOs-2 cells in T-152 cm<sup>2</sup> culture flasks in *growth media*.
  - Keep the density at 5-6 x 10<sup>5</sup> cells/ml
  - Passage cells 1:3 twice a week
- 20 • Add Trypsin EDTA (Gibco BRL Cat # 25300-020) and incubate
- Resuspend cells in plating media and transfer into growth media.

## Wash Media:

- HBSS Low Glucose Without Phenol Red (Gibco BRL Cat # 14175-095), 1% Ab-Am

## Plating Media:

- 25 • D-MEM Low Glucose Without Phenol Red (Gibco BRL Cat # 11054-020), 1% Ab-Am
- D-MEM

Stripped FBS (Hyclone Cat# SH30068.03 Lot # AHM9371 )

Ab-Am

## Transfection / Treatment Media:

- 30 • D-MEM Low Glucose Without Phenol Red only

T-152 cm<sup>2</sup> culture flask:

- Use Corning Coastr T-152 cm<sup>2</sup> culture flask (Cat # 430825) to grow the cells



-104-

Flat well Plates:

- Use well plate to plate cells
- Use Deep well plate sterile to make up treatment media.

5 Luciferase Assay Reagent:

- Use Steady-Glo Luciferase Reagent from Promega (Cat # E2550) Consists of:
  - a. E2533 Assay Substrate, lyophilized product and
  - b. E2543 Assay Buffer.
- Thaw at room temperature

10 • Store

DAY 1: Cell Plating:

Cell Harvesting

Aspirate media from culture flask, rinse cells with HBSS and aspirate.

Add trypsin and incubate.

15 When cells appear detached, resuspend cells in *growth media*.

Transfer into a new flask with fresh *growth media* for passaging the cells.

Plate well plates and two extra plates

A. Cell Count

Mix the cell suspension using pipette

20 Use *Hematocytometer* to count the cells

Load cell suspension onto the hemocytometer chamber

Count cells.

Plate seeding:

Use plating media 10 % Stripped FBS in D-MEM Low Glucose, Without Phenol Red, 1%

25 Ab-Am

Plate 14 plates @ 165  $\mu$ l / well.

In sterile flask add cell suspension

to *plating media*.

Mix .

30 Add cells / well.

Place the cells in the incubator.

Cells should be about 75 % confluent prior to transfection.

-105-

**Step 1: DNA and Media**

Add plain DMEM media to tubes for mixing the DNA

Add the Reporter gene pFR-LUC

- 5 Add the Gal4-RXR-DEF and VP16-VDR-LBD

**Step 2: FuGENE and Media**

Prepare plain DMEM media in a tubes for mixing FuGENE

Add *FuGENE 6 Transfection Reagent*

- 10 Incubate

**Step 3: FuGENE , DNA and Media Complex**

Add FuGENE Media complex from step 2 to DNA Media complex from step1

Incubate

15

**Step 4: FuGENE , DNA and Media Complex to-well plate**

Add FuGENE-DNA-Media complex from step 3 to each plate

Incubate.

20

**Day 3: Dosing****Treatment preparation**

Allow for transfection time

Make a stock solution of the compounds in DMSO

Vortex until all the compounds has been dissolved.

- 25 Further dilute in D-MEM (Low Glucose – With out Phenol Red)

Add compounds in quadruplicate to give final volume

Incubate.

**Day 4: Luciferase Assay**

Read the plates after drug treatment

- 30 Remove part of media from all the wells and leave remainder

Add Steady-Glo Luciferase Reagent mixture / wells

Incubate

-106-

Count each well using a Luminescence counter, Top Count NXT by Packard

Set a delay between plates to reduce the background.

(2) Materials and Method for The Caco-2 Cell Assay:

5 Caco-2 cells, grown in phenol red free, DMEM (Invitrogen, Carlsbad, CA)  
containing 10 % charcoal-stripped FCS (Hyclone, Logan, UT), were transfected  
with Fugene 6 reagent (Roche Diagnostics, Indianapolis, IN). Cells (5000/well)  
were plated 18 h before transfection in a 96 well plate. The Cells were transfected  
with Gal4-responsive reporter pFRLuc (150 ng, Stratagene, La Jolla CA) and the  
10 receptor expression vector pGal4-VDR-LBD (10 ng), along with Fugene 6  
reagent (0.2  $\mu$ l/well). The DNA-Fugene complex was formed by incubating the  
mixture for 30 min at room temperature. The cells were transfected in triplicate  
for 5 h, and treated with various concentrations of VDR ligands (from 0.01 nM to  
10,000 nM concentration range) 18h post-transfection. The luciferase activity  
15 was quantified using Steady-Glo reagent kit (Promega, Madison, WI) as per  
manufacturer's specifications.

(3) Materials and Method for The OCN Promoter Assay:

20 The activation of osteocalcin by VDR ligands was evaluated in a rat  
osteoblast-like cell line RG-15 (ROS 17/2.8) stably expressing rat osteocalcin  
promoter fused with luciferase reporter gene. The stable cell lines were  
established as reported before (Activation of Osteocalcin Transcription involves  
interaction of protein kinase A- and Protein kinase C-dependent pathways.  
25 Boguslawski, G., Hale, L. V., Yu, X.-P., Miles, R. R., Onyia, J. E., Santerre R.  
F., Chandrasekhar, S. J Biol. Chem. 275, 999-1006, 2000). Confluent RG-15  
cells maintained in DMEM/F-12 medium (3:1) containing 5% FBS, 300  $\mu$ g/ml  
G418 and at 37°C under 5% CO<sub>2</sub>/95% air atmosphere were trypsinized (0.25%  
trypsin) and plated into white opaque 96-well cell culture plates (25000  
30 cells/well). After 24 hr, cells (in DMEM/F-12 medium + 2% FBS) were treated  
with various concentrations of compounds, dissolved in DMSO. The final  
DMSO concentration remained at 0.01% (v/v). After 48 hr treatment, the

-107-

medium was removed, cells were lysed with 50  $\mu$ l of lysis buffer (From Luciferase reporter assay system, Roche Diagnostics, Indianapolis, IN) and assayed for luciferase activity using the Luciferase Reporter Gene Assay kit from Boehringer Mannheim as per manufacturer's specifications.

5

(4) Materials and Method for The Mouse Hypercalcemia Assay:

Weanling, virus -antibody-free, five to six weeks old female DBF mice (Harlan, Indianapolis, IN) are used for all the studies. Animals are allowed to acclimate to local vivarium conditions for 2 days. Mice are maintained on a 12 hr light/dark cycle at 22°C with ad lib access to food (TD 5001 with 1.2% Ca and 0.9%P, Teklad, Madison, WI) and water. The animals then are divided into groups with 4-5 mice per group. Different doses of test compounds prepared in 10% Ethanol and 90% sesame oil are administered to mice orally via gavage for 6 days.  $1\alpha$ -25(OH) $_2$ D $_3$  0.5 $\mu$ g/kg/d was also given to one group of mice as the positive control. Serum ionized calcium is evaluated at 6 hours after the last dosing under isoflurane anesthesia by Ciba-Corning Ca++/PH Analyzer, (Model 634, Chiron Diagnostics Corp., East Walpole, MA). Raw data of group differences is assessed by analysis of variance (ANOVA) using Fisher's protected least significant difference (PLSD) where the significance level was P< 0.05.

20 (5) The Keratinocyte Proliferation Assay:

KERtr cells (Human skin keratinocyte transformed with a retrovirus vector, obtained from ATCC) were plated in 96-well flat-bottomed plates (3000 cells/well) in 100  $\mu$ l keratinocyte serum free medium supplemented with bovine pituitary extract in the absence of EGF (Life Technologies, Rockville, MD) and incubated at 37°C for two days. The cells were treated with various concentrations of VDR ligands (ten-fold serial dilution from 10,000 nM to 0.1 nM in triplicate), dissolved in 100  $\mu$ l keratinocyte serum free medium supplemented with bovine pituitary extract in the absence of EGF and incubated at 37°C for 72hr. BrdU (5-bromo-2'-deoxyuridine) incorporation was analyzed as a measure of DNA replication (Cell proliferation ELISA kit, Roche Diagnostics, Indianapolis, IN) and absorbance was measured at 405 nm. Potency values (IC $_{50}$ ) values were determined as the concentration (nM) of compound that elicited a half-maximal response.

30

**(6) Materials and Method for human IL-10 Induction Assay:**

-109-

serum (Invitrogen, Carlsbad, CA), are plated at 5500 cell per well in a 96-well plate in a total volume of 100  $\mu$ l/well. The cells are kept in the 96-well plate for 6 days to differentiate them to small intestinal cells that express the calcium transporter, CaT1. On day 3 after plating, old media is removed and replaced with fresh media (150  $\mu$ l/well). On

5 day 6 the old media is removed and the cells are kept in treatment media (180  $\mu$ l/well) that contained 10 % charcoal stripped fetal bovine serum (Hyclone, Logan, UT) in DMEM (low glucose, without phenol red; Invitrogen, Carlsbad, CA). The cells are treated with various concentrations of VDR ligands (from 0.01 nM to 10,000 nM concentration range) prepared in treatment media (20  $\mu$ l/well). Twenty hours post-

10 treatment, total RNA is prepared by RNeasy 96 method as described by the manufacturer (Qiagen, Valencia, CA). The RNA is reverse transcribed and amplified for human CaT1 and GAPDH (control) messages by quantitative RT-PCR using ABI PRISM 7900HT Sequence Detection System according to manufacturer's instructions (Applied Biosystems, Foster City, CA). Optimized primer pairs and probes for human CaT1 and


15 GAPDH genes are obtained commercially (Applied Biosystems, Foster City, CA). Each 20  $\mu$ l quantitative RT-PCR reaction in a 384-well Taqman PCR plate consists of forward and reverse primers (900 nM), Taqman probe (200 nM), total RNA (4  $\mu$ l from each well of the 96-well culture plate) and 10  $\mu$ l of Taqman Universal PCR Master Mix (Roche Diagnostics, Indianapolis, IN). Reactions are incubated at 48°C for 30 minutes, followed

20 by 10 minutes at 95°C and subjected to 40 cycles of PCR (95°C for 15 seconds followed by 60°C for 1 minute). GAPDH is used as an internal control and its primer and probe set are obtained commercially (Applied Biosystems, Foster City, CA).

-110-

## WE CLAIM:

- 5 1. A compound represented by formula I or a pharmaceutically acceptable salt or a prodrug derivative thereof:-



wherein;




- R and R' are independently C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>1</sub>-C<sub>5</sub> fluoroalkyl, or together R and R' form a substituted or unsubstituted, saturated or unsaturated carbocyclic ring having from 3 to 8 carbon atoms;

R<sub>PH</sub> is hydrogen or methyl;





- R<sub>1</sub> and R<sub>2</sub> are independently selected from the group consisting of hydrogen, halo, C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>1</sub>-C<sub>5</sub> fluoroalkyl, -O-C<sub>1</sub>-C<sub>5</sub> alkyl, -S-C<sub>1</sub>-C<sub>5</sub> alkyl, -O-C<sub>1</sub>-C<sub>5</sub> fluoroalkyl, -CN, -NO<sub>2</sub>, acetyl, -S-C<sub>1</sub>-C<sub>5</sub> fluoroalkyl, C<sub>2</sub>-C<sub>5</sub> alkenyl, C<sub>3</sub>-C<sub>5</sub> cycloalkyl, and C<sub>3</sub>-C<sub>5</sub> cycloalkenyl;

L<sub>1</sub> and L<sub>2</sub> and L<sub>3</sub> are divalent linking groups independently selected from the group consisting of


a bond ,










-111-




$$\text{R40}$$

$$\text{R40}$$


$$\text{R40}$$




-112-



where m is 0, 1 or 2, X<sub>1</sub> is oxygen or sulfur, and each R<sub>40</sub> is independently hydrogen,

5 C<sub>1</sub>-C<sub>5</sub> alkyl, or C<sub>1</sub>-C<sub>5</sub> fluoroalkyl;

R<sub>BOH</sub> is

3-methyl-3-hydroxypentyl,  
 3-methyl-3-hydroxypentenyl,  
 3-methyl-3-hydroxypentynyl,  
 10 3-ethyl-3-hydroxypentyl,  
 3-ethyl-3-hydroxypentenyl,  
 3-ethyl-3-hydroxypentynyl,  
 3-ethyl-3-hydroxy-4-methylpentyl,  
 3-ethyl-3-hydroxy-4-methylpentenyl,  
 15 3-ethyl-3-hydroxy-4-methylpentynyl,  
 3-propyl-3-hydroxypentyl,  
 3-propyl-3-hydroxypentenyl,  
 3-propyl-3-hydroxypentynyl,  
 1-hydroxy-2-methyl-1-(methylethyl)propyl,  
 20 1-hydroxycyclopentenyl,  
 1-hydroxycyclohexenyl,  
 1-hydroxycycloheptenyl,  
 1-hydroxycyclooctenyl,  
 1-hydroxycyclopropyl,  
 25 1-hydroxycyclobutyl,  
 1-hydroxycyclopentyl,

-113-

1-hydroxycyclohexyl,  
 1-hydroxycycloheptyl, or  
 1-hydroxycyclooctyl;

provided, however, that when

5

$R_{BOH}$  is

10

3-methyl-3-hydroxypentyl,  
 3-methyl-3-hydroxypentenyl,  
 3-methyl-3-hydroxypentynyl,  
 3-ethyl-3-hydroxypentyl,  
 3-ethyl-3-hydroxypentenyl,  
 3-ethyl-3-hydroxypentynyl,  
 3-ethyl-3-hydroxy-4-methylpentyl,  
 3-ethyl-3-hydroxy-4-methylpentenyl,  
 3-ethyl-3-hydroxy-4-methylpentynyl,  
 3-propyl-3-hydroxypentyl,  
 3-propyl-3-hydroxypentenyl,  
 3-propyl-3-hydroxypentynyl, or  
 1-hydroxy-2-methyl-1-(methylethyl)propyl;

15

then  $L_1$  and  $L_2$  combine as a bond; and

20

$R_C$  is

25

$-CO_2H$ ,  
 $-CO_2Me$ ,  
 $-CO_2Et$ ,  
 $-C(O)CH_2S(O)Me$ ,  
 $-C(O)CH_2S(O)Et$ ,  
 $-C(O)CH_2S(O)_2Me$ ,  
 $-C(O)CH_2S(O)_2Et$ ,  
 $-C(O)CH_2CH_2S(O)Me$ ,  
 $-C(O)CH_2CH_2S(O)Et$ ,  
 $-C(O)CH_2CH_2S(O)_2Me$ ,  
 $-C(O)CH_2CH_2S(O)_2Et$ ,  
 $-C(O)CHMeCH_2CO_2H$

30

-114-

- 5 - -  
-C(O)C(O)OH,  
-C(O)C(O)NH<sub>2</sub>,  
-C(O)C(O)NHMe,  
-C(O)C(O)NMe<sub>2</sub>,  
-C(O)NH<sub>2</sub>, C(O)NMe<sub>2</sub>,  
-C(O)NHS(O)Me,  
-C(O)NHSO<sub>2</sub>Me,  
-C(O)-NH-5-tetrazolyl,  
-C(O)NMe-5-tetrazolyl,  
10 -C(O)NHS(O)Me,  
-C(O)NHS(O)Et,  
-C(O)NHSO<sub>2</sub>Me,  
-C(O)NHSO<sub>2</sub>Et,  
-C(O)NHS(O)iPr,  
15 -C(O)NHSO<sub>2</sub>iPr,  
-C(O)NHS(O)nPr,  
-C(O)NHSO<sub>2</sub>nPr,  
-C(O)NHCH<sub>2</sub>S(O)Me,  
-C(O)NHCH<sub>2</sub>S(O)Et,  
20 -C(O)NHCH<sub>2</sub>SO<sub>2</sub>Me,  
-C(O)NHCH<sub>2</sub>SO<sub>2</sub>Et,  
-C(O)NHCH<sub>2</sub>CH<sub>2</sub>S(O)Me,  
-C(O)NHCH<sub>2</sub>CH<sub>2</sub>S(O)Et,  
-C(O)NHCH<sub>2</sub>CH<sub>2</sub>SO<sub>2</sub>Me,  
25 -C(O)NHCH<sub>2</sub>CH<sub>2</sub>SO<sub>2</sub>Et,  
-C(O)NH<sub>2</sub>,  
-C(O)NMe<sub>2</sub>,  
-C(O)NH-CH<sub>2</sub>-C(O)OH,  
-C(O)NH-CH(Me)-C(O)OH,  
30 -C(O)NH-CH(F)-C(O)OH,  
-C(O)NH-CH(CF<sub>3</sub>)-C(O)OH,  
-C(O)NH-CH(OH)-C(O)OH,

-115-


- 5  
10  
15  
20  
25  
30
- C(O)NH-CH(cyclopropyl)-C(O)OH,
  - C(O)NH-C(Me)<sub>2</sub>-C(O)OH,
  - C(O)NH-C(Me)<sub>2</sub>-C(O)OH,
  - C(O)NH-CF(Me)-C(O)OH,
  - C(O)NH-C(Me)(CF<sub>3</sub>)-C(O)OH,
  - C(O)NH-C(Me)(OH)-C(O)OH,
  - C(O)NH-C(Me)(cyclopropyl)-C(O)OH,
  - C(O)NMe-CH<sub>2</sub>-C(O)OH,
  - C(O)NMe-CH(Me)-C(O)OH,
  - C(O)NMe-CH(F)-C(O)OH,
  - C(O)NMe-CH(CF<sub>3</sub>)-C(O)OH,
  - C(O)NMe-CH(OH)-C(O)OH,
  - C(O)NMe-CH(cyclopropyl)-C(O)OH,
  - C(O)NMe-C(Me)<sub>2</sub>-C(O)OH,
  - C(O)NMe-CF(Me)-C(O)OH,
  - C(O)NMe-C(Me)(CF<sub>3</sub>)-C(O)OH,
  - C(O)NMe-C(Me)(OH)-C(O)OH,
  - C(O)NMe-C(Me)(cyclopropyl)-C(O)OH,
  - CH<sub>2</sub>-CO<sub>2</sub>H,
  - CH<sub>2</sub>-5-tetrazolyl,
  - CH<sub>2</sub>CO<sub>2</sub>Me,
  - CH<sub>2</sub>CO<sub>2</sub>Et,
  - CH<sub>2</sub>NHS(O)Me,
  - CH<sub>2</sub>NHS(O)Et,
  - CH<sub>2</sub>NHSO<sub>2</sub>Me,
  - CH<sub>2</sub>NHSO<sub>2</sub>Et,
  - CH<sub>2</sub>NHS(O)iPr,
  - CH<sub>2</sub>NHSO<sub>2</sub>iPr,
  - CH<sub>2</sub>NHS(O)nPr,
  - CH<sub>2</sub>NHSO<sub>2</sub>nPr,
  - CH<sub>2</sub>NHCH<sub>2</sub>CH<sub>2</sub>SO<sub>2</sub>CH<sub>3</sub>,
  - CH<sub>2</sub>NH(CH<sub>2</sub>CO<sub>2</sub>H),

-116-









- 5                   -CH<sub>2</sub>N(C(O)Me)(CH<sub>2</sub>CO<sub>2</sub>H),  
                  -CH<sub>2</sub>-N-pyrrolidin-2-one,  
                  -CH<sub>2</sub>-(1-methylpyrrolidin-2-one-3-yl),  
                  -CH<sub>2</sub>S(O)Me,  
                  -CH<sub>2</sub>S(O)Et,  
                  -CH<sub>2</sub>S(O)<sub>2</sub>Me,  
                  -CH<sub>2</sub>S(O)<sub>2</sub>Et,  
                  -CH<sub>2</sub>S(O)iPr,  
                  -CH<sub>2</sub>S(O)<sub>2</sub>iPr,  
10                  -CH<sub>2</sub>S(O)nPr,  
                  -CH<sub>2</sub>S(O)<sub>2</sub>nPr,  
                  -CH<sub>2</sub>CO<sub>2</sub>H, CH<sub>2</sub>C(O)NH<sub>2</sub>,  
                  -CH<sub>2</sub>C(O)NMe<sub>2</sub>,  
                  -CH<sub>2</sub>C(O)NHMe,  
15                  -CH<sub>2</sub>C(O)-N-pyrrolidine,  
                  -CH<sub>2</sub>S(O)<sub>2</sub>Me,  
                  -CH<sub>2</sub>S(O)Me,  
                  -CH(OH)CO<sub>2</sub>H,  
                  -CH(OH)C(O)NH<sub>2</sub>,  
20                  -CH(OH)C(O)NHMe,  
                  -CH(OH)C(O)NMe<sub>2</sub>,  
                  -CH(OH)C(O)NEt<sub>2</sub>,  
                  -CH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub>H,  
                  -CH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub>Me,  
25                  -CH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub>Et,  
                  -CH<sub>2</sub>CH<sub>2</sub>C(O)NH<sub>2</sub>,  
                  -CH<sub>2</sub>CH<sub>2</sub>C(O)NHMe,  
                  -CH<sub>2</sub>CH<sub>2</sub>C(O)NMe<sub>2</sub>,  
                  -CH<sub>2</sub>CH<sub>2</sub>-5-tetrazolyl,  
30                  -CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>Me,  
                  -CH<sub>2</sub>CH<sub>2</sub>S(O)Me,  
                  -CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>Et,

-117-





- 5  
10  
15  
20
- CH<sub>2</sub>CH<sub>2</sub>S(O) Et,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)iPr,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>iPr,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)nPr,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>nPr,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)NH<sub>2</sub>,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)NHMe,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)NMe<sub>2</sub>,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>NH<sub>2</sub>,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>NHMe,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>NMe<sub>2</sub>,
  - CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>S(O)Me,
  - CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>S(O)Et,
  - CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>Me,
  - CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>Et,
  - CH(Me)CH<sub>2</sub>C(O)OH,
  - C(Me)<sub>2</sub>CH<sub>2</sub>C(O)OH,
  - SO<sub>3</sub>H,
  - 5-tetrazolyl,




-118-



-119-



or



-1,3,4-oxadiazolin-2-one-5-yl,

-imidazolidine-2,4-dione-5-yl,

-1,3-thiazolidine-2,4-dione-5-methylidene,

-isoxazol-3-ol-yl, or

-1,3,4-oxadiazolin-2-thione-5-yl.









-120-

2. A compound according to Claim 1 or a pharmaceutically acceptable salt or ester prodrug derivative thereof wherein

$R_{PH}$  is hydrogen;


$L_3$  are divalent linking groups independently selected from the group consisting of

a bond





R40


R40



R40



-121-



5

 $R_C$  is

10

15

$-\text{CO}_2\text{H},$   
 $-\text{CO}_2\text{Me},$   
 $-\text{CO}_2\text{Et},$   
 $-\text{C(O)CH}_2\text{S(O)Me},$   
 $-\text{C(O)CH}_2\text{S(O)Et},$   
 $-\text{C(O)CH}_2\text{S(O)}_2\text{Me},$   
 $-\text{C(O)CH}_2\text{S(O)}_2\text{Et},$   
 $-\text{C(O)CH}_2\text{CH}_2\text{S(O)Me},$   
 $-\text{C(O)CH}_2\text{CH}_2\text{S(O)Et},$

-122-

- 5                   -C(O)CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>Me,  
                  -C(O)CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>Et,  
                  -C(O)CHMeCH<sub>2</sub>CO<sub>2</sub>H  
                  -C(O)C(O)OH,  
                  -C(O)C(O)NH<sub>2</sub>,  
                  -C(O)C(O)NHMe,  
                  -C(O)C(O)NMe<sub>2</sub>,  
                  -C(O)NH<sub>2</sub>, C(O)NMe<sub>2</sub>,  
10                  -C(O)NHS(O)Me,  
                  -C(O)NHSO<sub>2</sub>Me,  
                  -C(O)-NH-5-tetrazolyl,  
                  -C(O)NMe-5-tetrazolyl,  
                  -C(O)NHS(O)Me,  
                  -C(O)NHS(O)Et,  
15                  -C(O)NHSO<sub>2</sub>Me,  
                  -C(O)NHSO<sub>2</sub>Et,  
                  -C(O)NHS(O)iPr,  
                  -C(O)NHSO<sub>2</sub>iPr,  
                  -C(O)NHS(O)nPr,  
20                  -C(O)NHSO<sub>2</sub>nPr,  
                  -C(O)NHCH<sub>2</sub>S(O)Me,  
                  -C(O)NHCH<sub>2</sub>S(O)Et,  
                  -C(O)NHCH<sub>2</sub>SO<sub>2</sub>Me,  
                  -C(O)NHCH<sub>2</sub>SO<sub>2</sub>Et,  
25                  -C(O)NHCH<sub>2</sub>CH<sub>2</sub>S(O)Me,  
                  -C(O)NHCH<sub>2</sub>CH<sub>2</sub>S(O)Et,  
                  -C(O)NHCH<sub>2</sub>CH<sub>2</sub>SO<sub>2</sub>Me,  
                  -C(O)NHCH<sub>2</sub>CH<sub>2</sub>SO<sub>2</sub>Et,  
                  -C(O)NH<sub>2</sub>,  
30                  -C(O)NMe<sub>2</sub>,  
                  -C(O)NH-CH<sub>2</sub>-C(O)OH,  
                  -C(O)NH-CH(Me)-C(O)OH,

-123-

- 5
- 10
- 15
- 20
- C(O)NH-CH(F)-C(O)OH,
  - C(O)NH-CH(CF<sub>3</sub>)-C(O)OH,
  - C(O)NH-CH(OH)-C(O)OH,
  - C(O)NH-CH(cyclopropyl)-C(O)OH,
  - C(O)NH-C(Me)<sub>2</sub>-C(O)OH,
  - C(O)NH-C(Me)<sub>2</sub>-C(O)OH,
  - C(O)NH-CF(Me)-C(O)OH,
  - C(O)NH-C(Me)(CF<sub>3</sub>)-C(O)OH,
  - C(O)NH-C(Me)(OH)-C(O)OH,
  - C(O)NH-C(Me)(cyclopropyl)-C(O)OH,
  - C(O)NMe-CH<sub>2</sub>-C(O)OH,
  - C(O)NMe-CH(Me)-C(O)OH,
  - C(O)NMe-CH(F)-C(O)OH,
  - C(O)NMe-CH(CF<sub>3</sub>)-C(O)OH,
  - C(O)NMe-CH(OH)-C(O)OH,
  - C(O)NMe-CH(cyclopropyl)-C(O)OH,
  - C(O)NMe-C(Me)<sub>2</sub>-C(O)OH,
  - C(O)NMe-CF(Me)-C(O)OH,
  - C(O)NMe-C(Me)(CF<sub>3</sub>)-C(O)OH,
  - C(O)NMe-C(Me)(OH)-C(O)OH,
  - C(O)NMe-C(Me)(cyclopropyl)-C(O)OH,

-124-

- 5
- CH<sub>2</sub>-CO<sub>2</sub>H,  
-CH<sub>2</sub>-5-tetrazolyl,  
-CH<sub>2</sub> CO<sub>2</sub>Me,  
-CH<sub>2</sub>CO<sub>2</sub>Et,  
-CH<sub>2</sub>NHS(O)Me,  
-CH<sub>2</sub>NHS(O)Et,  
-CH<sub>2</sub>NHSO<sub>2</sub>Me,  
-CH<sub>2</sub>NHSO<sub>2</sub>Et,  
-CH<sub>2</sub>NHS(O)iPr,  
10 -CH<sub>2</sub>NHSO<sub>2</sub>iPr,  
-CH<sub>2</sub>NHS(O)nPr,  
-CH<sub>2</sub>NHSO<sub>2</sub>nPr,  
-CH<sub>2</sub>NHCH<sub>2</sub>CH<sub>2</sub>SO<sub>2</sub>CH<sub>3</sub>,  
-CH<sub>2</sub>NH(CH<sub>2</sub>CO<sub>2</sub>H),  
15 -CH<sub>2</sub>N(C(O)Me)(CH<sub>2</sub>CO<sub>2</sub>H),  
-CH<sub>2</sub>-N-pyrrolidin-2-one,  
-CH<sub>2</sub>-(1-methylpyrrolidin-2-one-3-yl),  
-CH<sub>2</sub>S(O)Me,  
-CH<sub>2</sub>S(O)Et,  
20 -CH<sub>2</sub>S(O)<sub>2</sub>Me,  
-CH<sub>2</sub>S(O)<sub>2</sub>Et,  
-CH<sub>2</sub>S(O)iPr,  
-CH<sub>2</sub>S(O)<sub>2</sub>iPr,  
-CH<sub>2</sub>S(O)nPr,  
25 -CH<sub>2</sub>S(O)<sub>2</sub>nPr,  
-CH<sub>2</sub>CO<sub>2</sub>H, CH<sub>2</sub>C(O)NH<sub>2</sub>,  
-CH<sub>2</sub>C(O)NMe<sub>2</sub>,  
-CH<sub>2</sub>C(O)NHMe,  
-CH<sub>2</sub>C(O)-N-pyrrolidine,  
30 -CH<sub>2</sub>S(O)<sub>2</sub>Me,  
-CH<sub>2</sub>S(O)Me,  
-CH(OH) CO<sub>2</sub>H,

-125-

- 5
- CH(OH)C(O)NH<sub>2</sub>,
  - CH(OH)C(O)NHMe,
  - CH(OH)C(O)NMe<sub>2</sub>,
  - CH(OH)C(O)NEt<sub>2</sub>,
  - CH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub>H,
  - CH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub>Me,
  - CH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub>Et,
  - CH<sub>2</sub>CH<sub>2</sub>C(O)NH<sub>2</sub>,
  - CH<sub>2</sub>CH<sub>2</sub>C(O)NHMe,

10

  - CH<sub>2</sub>CH<sub>2</sub>C(O)NMe<sub>2</sub>,
  - CH<sub>2</sub>CH<sub>2</sub>-5-tetrazolyl,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>Me,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)Me,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>Et,

15

  - CH<sub>2</sub>CH<sub>2</sub>S(O) Et,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)iPr,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>iPr,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)nPr,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>nPr,

20

  - CH<sub>2</sub>CH<sub>2</sub>S(O)NH<sub>2</sub>,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)NHMe,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)NMe<sub>2</sub>,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>NH<sub>2</sub>,
  - CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>NHMe,

25










  - CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>NMe<sub>2</sub>,
  - CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>S(O)Me,
  - CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>S(O)Et,
  - CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>Me,
  - CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>Et,

30





  - SO<sub>3</sub>H,

-126-


-5-tetrazolyl,



-127-



or



-1,3,4-oxadiazolin-2-one-5-yl,

-imidazolidine-2,4-dione-5-yl,

-1,3-thiazolidine-2,4-dione-5-methylidene,


-isoxazol-3-ol-yl, or

-1,3,4-oxadiazolin-2-thione-5-yl.



-128-








3. A compound represented by formula (II) or a pharmaceutically acceptable salt or an ester prodrug derivative thereof:



wherein;

- 5      R and R' are independently methyl or ethyl;  
       R1 and R2 are independently hydrogen, halo, -CF<sub>3</sub>, methyl, ethyl, or cyclopropyl;  
       L<sub>1</sub> and L<sub>2</sub> are independently divalent linking groups independently selected from

a bond



where m is 0 or 1;

R<sub>BOH</sub> is selected from

- 15      1-hydroxycyclopentenyl,  
       1-hydroxycyclohexenyl,  
       1-hydroxycyclopentyl, or  
       1-hydroxycyclohexyl, and


R<sub>C</sub> is a group selected from

-129-

- 5
- 10
- 15
- 20
- 25
- 30
- CO<sub>2</sub>H,
  - CO<sub>2</sub>Me,
  - CO<sub>2</sub>Et,
  - C(O)NH<sub>2</sub>,
  - C(O)NMe<sub>2</sub>,
  - C(O)NH-CH<sub>2</sub>-C(O)OH,
  - C(O)NH-CH(Me)-C(O)OH,
  - C(O)NH-CH(F)-C(O)OH,
  - C(O)NH-CH(CF<sub>3</sub>)-C(O)OH,
  - C(O)NH-CH(OH)-C(O)OH,
  - C(O)NH-CH(cyclopropyl)-C(O)OH,
  - C(O)NH-C(Me)<sub>2</sub>-C(O)OH,
  - C(O)NH-C(Me)<sub>2</sub>-C(O)OH,
  - C(O)NH-CF(Me)-C(O)OH,
  - C(O)NH-C(Me)(CF<sub>3</sub>)-C(O)OH,
  - C(O)NH-C(Me)(OH)-C(O)OH,
  - C(O)NH-C(Me)(cyclopropyl)-C(O)OH,
  - C(O)NMe-CH<sub>2</sub>-C(O)OH,
  - C(O)NMe-CH(Me)-C(O)OH,
  - C(O)NMe-CH(F)-C(O)OH,
  - C(O)NMe-CH(CF<sub>3</sub>)-C(O)OH,
  - C(O)NMe-CH(OH)-C(O)OH,
  - C(O)NMe-CH(cyclopropyl)-C(O)OH,
  - C(O)NMe-C(Me)<sub>2</sub>-C(O)OH,
  - C(O)NMe-CF(Me)-C(O)OH,
  - C(O)NMe-C(Me)(CF<sub>3</sub>)-C(O)OH,
  - C(O)NMe-C(Me)(OH)-C(O)OH,
  - C(O)NMe-5-tetrazolyl,
  - C(O)NMe-C(Me)(cyclopropyl)-C(O)OH, or
  - C(O)-NH-5-tetrazolyl.

-130-

4. A compound represented by formula (III) or a pharmaceutically acceptable salt or an ester prodrug derivative thereof:



wherein;

5

R and R' are independently methyl or ethyl;

R1 and R2 are independently hydrogen, halo, -CF<sub>3</sub>, methyl, ethyl, or cyclopropyl;

R<sub>BOH</sub> is selected from

10

3-methyl-3-hydroxypentyl,

3-methyl-3-hydroxypentenyl,

3-methyl-3-hydroxypentynyl,

3-ethyl-3-hydroxypentyl,

3-ethyl-3-hydroxypentenyl,

3-ethyl-3-hydroxypentynyl,

3-propyl-3-hydroxypentyl,

15

3-propyl-3-hydroxypentenyl,

3-propyl-3-hydroxypentynyl,

3-ethyl-3-hydroxy-4-methylpentyl,

3-ethyl-3-hydroxy-4-methylpentenyl,

3-ethyl-3-hydroxy-4-methylpentynyl, or

20

1-hydroxy-2-methyl-1-(methylethyl)propyl;

and

R<sub>C</sub> is a group selected from

25

-CO<sub>2</sub>H,

-CO<sub>2</sub>Me,

-CO<sub>2</sub>Et,

-C(O)NH<sub>2</sub>,

-C(O)NMe<sub>2</sub>,

-C(O)NH-CH<sub>2</sub>-C(O)OH,

-C(O)NH-CH(Me)-C(O)OH,

-131-

- 5
- C(O)NH-CH(F)-C(O)OH,
  - C(O)NH-CH(CF<sub>3</sub>)-C(O)OH,
  - C(O)NH-CH(OH)-C(O)OH,
  - C(O)NH-CH(cyclopropyl)-C(O)OH,
  - C(O)NH-C(Me)<sub>2</sub>-C(O)OH,
  - C(O)NH-C(Me)<sub>2</sub>-C(O)OH,
  - C(O)NH-CF(Me)-C(O)OH,
  - C(O)NH-C(Me)(CF<sub>3</sub>)-C(O)OH,
  - C(O)NH-C(Me)(OH)-C(O)OH,

10

  - C(O)NH-C(Me)(cyclopropyl)-C(O)OH,
  - C(O)NMe-CH<sub>2</sub>-C(O)OH,
  - C(O)NMe-CH(Me)-C(O)OH,
  - C(O)NMe-CH(F)-C(O)OH,
  - C(O)NMe-CH(CF<sub>3</sub>)-C(O)OH,
  - C(O)NMe-CH(OH)-C(O)OH,
  - C(O)NMe-CH(cyclopropyl)-C(O)OH,
  - C(O)NMe-C(Me)<sub>2</sub>-C(O)OH,
  - C(O)NMe-CF(Me)-C(O)OH,
  - C(O)NMe-C(Me)(CF<sub>3</sub>)-C(O)OH,

15


  - C(O)NMe-C(Me)(OH)-C(O)OH,
  - C(O)NMe-5-tetrazolyl,
  - C(O)NMe-C(Me)(cyclopropyl)-C(O)OH, or
  - C(O)-NH-5-tetrazolyl.

20


- 25
5. The compound represented by formula (AA-1) to (AA-33) or a pharmaceutically acceptable salt or prodrug derivative thereof:

AA-1)


-132-



AA-2)




AA-3)




5

AA-4)




AA-5)


-133-




AA-6)



AA-7)




AA-8)




AA-9)


-134-




AA-10)



AA-11)




AA-12)




AA-13)


-135-




AA-14)



AA-15)




AA-16)




AA-17)




-136-




AA-18)



AA-19)




AA-20)




AA-21)


-137-




AA-22)



AA-23)




AA-24)




AA-25)


-138-




AA-26)



AA-27)




AA-28)




AA-29)


-139-




AA-30)



AA-31)




AA-32)



AA-33)

or


-140-




6. The compound represented by formula (BB-1) to (BB-3) or a pharmaceutically acceptable salt or prodrug derivative thereof:

5


BB-1)



BB-2)




BB-3)




-141-

BB-4)




BB-5)




5

BB-6)




BB-7)




BB-8)


-142-




BB-9)



BB-10)




BB-11)




BB-12)


-143-




BB-13)



BB-14)




BB-15)




BB-16)




-144-




BB-17)



BB-18)




BB-19)




BB-20)


-145-




BB-21)



BB-22)




BB-23)




BB-24)

-146-




BB-25)




5

BB-26)




BB-27)




BB-28)


-147-




BB-29)



BB-30)




BB-31)




BB-32)

-148-



, or


BB-33)




5

7. The compound represented by formula (CC-1) to (CC-44) or a pharmaceutically acceptable salt or prodrug derivative thereof:

CC-1)




CC-2)




-149-

CC-3)




CC-4)




5

CC-5)




CC-6)




CC-7)


-150-




CC-8)



CC-9)




CC-10)




CC-11)


-151-




CC-12)



CC-13)




CC-14)




CC-15)



-152-




CC-16)




5

CC-17)




CC-18)




-153-

CC-19)




CC-20)




5

CC-21)




CC-22)




CC-23)


-154-




CC-24)



CC-25)




CC-26)




CC-27)


-155-



CC-28)




CC-29)




5

CC-30)




CC-31)


-156-




CC-32)



CC-33)




CC-34)




CC-35)

-157-




CC-36)




CC-37)

5.




CC-38)




CC-39)


-158-




CC-40)



CC-41)




CC-42)




CC-43)

-159-




, or

CC-44)



5

8. The compound represented by the formula:




;

or a pharmaceutically acceptable salt or prodrug derivative thereof.

10

9. The compound represented by the formula:



;

or a pharmaceutically acceptable salt or prodrug derivative thereof.



-160-

10. The prodrug derivative of the compound of claim 1 to 9 wherein the prodrug is a methyl ester; ethyl ester; N,N-diethylglycolamido ester; or morpholinylethyl ester.

11. The salt derivative of the compound of claim 1 to 9 wherein the salt is sodium  
5 or potassium.

12. A pharmaceutical formulation comprising the compound of claim 1 to 9 together with a pharmaceutically acceptable carrier or diluent.

10 13. A formulation for treating osteoporosis comprising:  
Ingredient (A1): the vitamin D receptor modulator of claim 1,  
represented by formula (I);

Ingredient (B1):

one or more co-agents selected from the group consisting of:

- 15 a. estrogens,  
b. androgens,  
c. calcium supplements,  
d. vitamin D metabolites,  
e. thiazide diuretics,  
20 f. calcitonin,  
g. bisphosphonates,  
h. SERMS, and  
i. fluorides; and

Ingredient (C1): optionally, a carrier or diluent.

25

14. The formulation of claim 13 wherein the weight ratio of (A1) to (B1) is from 10:1 to 1:1000.

15. A formulation for treating osteoporosis comprising:

30 Ingredient (A2): the vitamin D receptor modulator of claim 1  
represented by formula (I);

Ingredient (B2):

-161-

one or more co-agents that are conventional for treatment  
osteoporosis selected from the group consisting of:

- a. topical glucocorticoids ,
- b. salicylic acid,
- c. crude coal tar; and

Ingredient (C2): optionally, a carrier or diluent.

16. The formulation of claim 15 wherein the weight ratio of (A2) to (B2) is  
from 1:10 to 1:100000.

17. A method of treating a mammal to prevent or alleviate the pathological  
effects of Acne, Actinic keratosis, Alopecia , Alzheimer's disease, Bone maintenance in  
zero gravity, Bone fracture healing, Breast cancer, Chemoprevention of Cancer, Crohn's  
disease, Colon cancer, Type I diabetes, Host-graft rejection, Hypercalcemia , Type II  
diabetes, Leukemia, Multiple sclerosis, Myelodysplastic syndrome, Insufficient sebum  
secretion, Osteomalacia, Osteoporosis, Insufficient dermal firmness, Insufficient dermal  
hydration, Psoriatic arthritis, Prostate cancer, Psoriasis, Renal osteodystrophy,  
Rheumatoid arthritis, Scleroderma, Skin cancer, Systemic lupus erythematosus, Skin cell  
protection from , Mustard vesicants, Ulcerative colitis, Vitiligo, or Wrinkles; wherein the  
method comprises administering a pharmaceutically effective amount of at least one  
compound of claim 1 or 9.

18. The method of claim 17 for the treatment of psoriasis.

19. The method of claim 17 for the treatment of osteoporosis.

20. A method of claim 17 for treating a mammal to prevent or alleviate skin cell  
protection from Mustard vesicants.

21. A method of of treating a mammal to prevent or alleviate the pathological  
effects of Benign prostatic hyperplasia or bladder cancer wherein the method comprises  
administering a pharmaceutically effective amount of at least one compound of claim 1 or

-162-

9.

22. A method of treating or preventing disease states mediated by the Vitamin D receptor, wherein a mammal in need thereof is administered a pharmaceutically effective amount of the compound of Claim 1 to 9.

23. A compound as claimed in any one of Claims 1 to 9 for use in treating a mammal to prevent or alleviate the pathological effects of Acne, Actinic keratosis, Alopecia, Alzheimer's disease, Bone maintenance in zero gravity, Bone fracture healing, Breast cancer, Chemoprevention of Cancer, Crohn's disease, Colon cancer, Type I diabetes, Host-graft rejection, Hypercalcemia, Type II diabetes, Leukemia, Multiple sclerosis, Myelodysplastic syndrome, Insufficient sebum secretion, Osteomalacia, Osteoporosis, Insufficient dermal firmness, Insufficient dermal hydration, Psoriatic arthritis, Prostate cancer, Psoriasis, Renal osteodystrophy, Rheumatoid arthritis, Scleroderma, Skin cancer, Systemic lupus erythematosus, Skin cell protection from, Mustard vesicants, Ulcerative colitis, Vitiligo, or Wrinkles.

24. A compound as claimed in any one of Claims 1 to 9 for use in treating or preventing disease states mediated by the Vitamin D receptor.

20

25. A compound as claimed in any one of Claims 1 to 9 for use in treating a mammal to prevent or alleviate the pathological effects of Benign prostatic hyperplasia or bladder cancer.

26. A compound as claimed in Claim 1 substantially as hereinbefore described with reference to any of the Examples.

27. A process for preparing a compound as claimed in claim 1 substantially as hereinbefore described with reference to any of the Examples.

30

-163-

28. The use of a compound as claimed in claim 1 substantially as herein described with reference to any of the Assays and Tables for mediating the Vitamin D<sub>3</sub> receptor.